

## DIRECTIONS

Nutro 7—SEVIN INSECT KILLER is a wettable powder especially designed for spray application. It is highly effective against Japanese Beetles and also controls certain other insects. It may be used with most commonly used insecticides and fungicides. Mix at the rate of 2 level tablespoons per gallon of water. Apply with hose end or spray type pressure sprayer. For best results, add enough water to make a thin paste, stir thoroughly. Then add balance of water required. Agitate frequently while spraying. Apply when insects first appear and continue at weekly intervals.

## INSECTS CONTROLLED

**Shade Trees**—Bagworm, Birch Leafminer, Boxelder bug, Elm leaf aphid, Elm leaf beetle, Gypsy moth, lace bug, Mimosa webworm, Oak leaf miners, periodical cicada, psyllids, puss caterpillars, scale insects, tent caterpillars, thorn bugs, willow leaf beetles.

**Ornamental Shrubs and Flowers**—Apple Aphid, blister beetles, boxwood leaf miner, flea beetles, Japanese beetle, June beetle, leafhoppers, leaf rollers, mealy bugs, orange tortrix, plant bugs, rose aphid, rose slug, thrips (exposed).

**Lawn Insects**—Ants, chinch bugs, earwigs, leafhoppers, armyworm, fleas, millipeds, mosquitos, sod webworms or other lawn moths.

**Apples and Pears**—Use 2 tablespoons per gallon for codling moth, pear psylla, apple maggot, plum curculio, fruit tree leaf rollers, Japanese beetles, apple aphids, and San Jose scale. Do not apply before second cover spray to avoid fruit thinning. Apply at 10-day intervals in July for apple maggot. May be used up to 1 day before picking.

**Peaches and Apricots**—Use 1 tablespoonful per gallon for Oriental fruit fly, plum curculio, peach twig borer, cat facing bugs, San Jose scale, Japanese beetle. May be used 1 day before harvest on peaches; 3 days on apricots.

**Grapes**—Use 1 tablespoonful per gallon for leafhoppers, grape berry moth, Japanese beetles. May be applied up to harvest.

**Strawberries**—Use 3 tablespoons per gallon up to 1 day before picking for leaf roller and meadow spittlebug control. Make 3 to 5 post bloom summer sprays if needed.

**Beans**—Use 2 tablespoons per gallon. Spray until all foliage wet. Repeat 7 to 10 days. For Mexican bean beetles, leafhoppers, flea beetles, stink bugs, lima bean pod borer. May be used up to harvest. Do not feed vines to dairy or meat animals.

**Corn**—Use 2 tablespoons per gallon for corn earworm, corn borer, Fall armyworm, and grasshoppers. Apply up to day of harvest. Repeat at 2 to 3 day intervals.

**Cucumbers, Melons, and Squash**—Use 2 tablespoons per gallon. Apply as full cover spray for squash bugs, melon worms, cucumber beetles, and pickleworms.

**Potatoes**—Use 2 tablespoons per gallon for Colorado potato beetles, leafhoppers, flea beetles.

**Tomatoes, Peppers, Eggplants**—Use 2 tablespoons per gallon for tomato fruitworm, flea beetles, Fall army worms, stink bugs and leafhoppers. Apply up to day of picking.

## CAUTION:


HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid breathing of spray. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Avoid contamination of feed and foodstuffs. Note for physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Wash thoroughly after using. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

Rinse empty container thoroughly with water and discard it.


NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but Seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty, express or implied except as herein stated. Seller's liability is limited to the purchase price of the product and this shall be the buyers only remedy. Buyer assumes the risk of use of this product contrary to label instructions carelessly, or under conditions not intended by or not reasonably foreseeable to Seller.

E.P.A. Reg. No. 4185 302

SMITH-DO' GLASS BORDEN CHEMICAL  
DIVISION OF BORDEN INC  
NORFOLK, VIRGINIA 23501




Nutro



Sevin

Insect  
Killer



SE

Insect


Japanese Beetles, Boxelder Bugs, Leaf Beetles, Flea Beetles, Leaf Rollers, Periodical Cicada, Caterpillars, Thrips (exposed) and other insects.

ACTIVE INGREDIENT:  
Carbaryl (1-Naphthyl-N-methylcarbamate)  
INERT INGREDIENTS:

\*Sevin — T.M. Union Carbide

NET

CAUTION: KEEP OUT OF REACH OF CHILDREN  
See back panel for



*Nutro.*

TESTED

1975

Insecticide  
Fentioide  
pesticide

-302

**Insect  
Killer**



*Nutro.*

**SEVIN**

**Insect Killer**

Japanese Beetles, Bagworms, Aphids, Blister Beetles, Boxelder Bugs, Certain Leafminers, Elm Leaf Beetles, Flea Beetles, Leafhoppers, Leaf Rollers, Periodical Cicada, Scale Insects, Tent Caterpillars, Thrips (exposed), and many other insects.


ACTIVE INGREDIENT:  
Carbaryl (1-Naphthyl-N-methylcarbamate)\*  
INERT INGREDIENTS:

50%  
50%  
100%

\*Sevin — T.M. Union Carbide Chemicals Co.

**NET WT. 6 OZ.**

**CAUTION:** KEEP OUT OF REACH OF CHILDREN AND PETS.  
See back panel for additional cautions.



*Nutro.*

**Sevin**

**Insect  
Killer**

ACCEPTED WITH COMMENTS