8MITH-DOUGLASS DUSTING SULFUR (93%)

Fineness: Not less than 93% passing 325 mesh.

CAUTION

Sulfur may cause severe fruit and leaf injury to sulfur-sensitive crops. Do not use on Apricots, Anjou and Comice Pears, Cranberries, Cucurbits (Cucumbers, squash and some varieties of melons), Filberts, Spinach, Tung Trees, Walnuts and other sensitive plants.

Sulfur may burn foliage and fruit during periods of high temperatures.

exceed or are likely to exceed 90 F.

Do not use sulfur with oil sprays or within 3 to 4 weeks of an oil spray.

If the crop to be treated will be used for canning or freezing purposes, consult food processor before applying sulfur.

DIRECTIONS FOR USE

Smith-Douglass Dusting Sulfur is formulated for use in dust mixtures or alone as dust when a 325-mesh, free-flowing dust is required.

IMPORTANT: For best results apply when air is calm. Dust often enough to keep foliage and fruit covered with a light, uniform coating. Consult State Agricultural Experiment Stations for additional information, as the timing, number and rate of applications needed will vary with local conditions.

RECOMMENDATIONS

Fruit Crops

APPLES: Scab, powdery mildew - Apply 30 - 40 lb. per acre in the green tip to pink bud stage. Use on sulfur tolerant varieties only.

CHERRY: Blossom blight, brown rot, powdery mildew - Apply 50 lb. per acre in the popcorn, full bloom and petal fall stages; leaf spot, powdery mildew - Apply 50 lb. per lb. per acre at petal fall, shuck-fall and 2 weeks after shuck-fall.

GRAPES: (Vinifera Varieties): Rust, Powdery Mildew - Apply 15 to 20 lb. per acre in 4 applications, when shoots are 6" long; shoots 12-15" long; 14 days afterward; 10-14 days thereafter to harvest. Caution: Sulfur is extremely phytotoxic to foliage of Concord and most other labrusca type grapes and should not be used on these varieties.

PEACH, NECTARINE: Brown rot, coryneum blight, rust, peach silver mite, powdery mildew - Apply 50 to 60 lb. per acre from popcorn stage to harvest.