


Front Panel

Bag Print
NUTRO CHINCH BUG CONTROL
Turf Food 20-5-5
Size: 15 Lbs.
March 22, 1968



BORDEN
CHEMICAL

NUTRO
(R)
CHINCH BUG CONTROL
Turf Food

TREATS & FEEDS
2,500 SQ. FT.
\$5.95

for exclusive use on lawns

CONTROLS CHINCH BUGS & BERMUDA MITES
Feeds grass back to good health

USDA Reg. No. 4185-359

YELLOW

ACTIVE INGREDIENTS:

*Carbophenothion S-[(p-Chlorophenylthio) methyl] 0,0-diethyl
phosphorodithioate

INERT INGREDIENTS:

TOTAL

3.15%
96.85%
100.00%

*Trithion - Reg. TM of Stauffer Chemical Co.
U. S. Pat. No. 2,793,224

**Equivalent to 8.22 Lbs. of Trithion actual per acre

NET WT.

15 LBS.

TREATS & FEEDS
2,500 SQ. FT.
\$5.95

NUTRO CHINCH BUG CONTROL Turf Food
(R)

YELLOW

CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS (See back panel for additional
cautions)

Back Panel

BORDEN
CHEMICAL
NUTRO
(R)
CHINCH BUG CONTROL
Turf Food

NET WT.

20 - 5 - 5

15 LBS.

Bag Print
NUTRO CHINCH BUG CONTROL
Turf Food 20-5-5
Size: 15 lbs.

Back Panel (continued)

followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. The Borden Chemical Co. and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks.

WARRANTY AND LIMITATIONS OF DAMAGES. The Borden Chemical Co. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale; subject to the risks referred to therein. The Borden Chemical Co. makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of the Borden Chemical Co. is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS FOR USE ON LAWN AREAS ONLY

YELLOW

Controls Chinch Bugs and Bermuda Mites

WHEN TO APPLY: Use immediately upon evidence that chinch bugs are at work. As a preventative, use as soon as weather is consistently above 80 degrees and repeat application at two month intervals, depending on length of chinch bug season in your area.

HOW TO APPLY: Use a fertilizer spreader and apply this bag to 2,500 sq. ft. (50' x 50') of lawn. Settings below are approximate for applying contents of this bag at suggested rate of 6 pounds per 1,000 sq. ft. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed.

Immediately after application, drench area with water or sprinkle thoroughly. Do not allow persons or pets on treated lawn until grass is completely dry.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION - HAZARDOUS IF SWALLOWED. Avoid breathing dust. Wear a dust respirator. Wear gloves. Avoid contact with skin and eyes. Wash hands with soap and water before eating, smoking, and after using. Wash all contaminated clothing with soap and hot water before re-use. Do not apply when weather conditions favor drift of dust from areas treated. Do not contaminate feed and foodstuffs.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed.

Keep children and pets off treated area until treated areas are well watered and completely dried.

STORE OUT OF REACH OF CHILDREN. Trithion, upon repeated, prolonged and careless use, may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL.

DESTROY THIS BAG BY BURNING WHEN EMPTY. Avoid exposure of persons or animals to smoke.

YELLOW

4-25-68


4185-359

Back Panel (continued)

Bag Print
NUTRO CHINCH BUG CONTROL
Turf Food 20-5-5
Size: 15 lbs.

GUARANTEED ANALYSIS

TOTAL NITROGEN (%)	20.00%
1.0% Ammoniacal Nitrogen	
13.5% Urea Nitrogen	
5.5% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	5.00%
WATER SOLUBLE POTASH (K ₂ O)	5.00%
Total Available Primary Plant Food - 30.00	
Chlorine (Cl)	1.00%
Sulphur (S)	1.00%
Iron (Fe)	0.2%



Derived from - Primary Plant Foods: Anhydrous Ammonia, Urea, Urea-Formaldehyde, Phosphoric Acid, Muriate of Potash and Sulphate of Potash.

Derived from - Secondary Plant Foods: Sulphate of Potash and Iron Phosphate.

Potential acidity equivalent to 800 lbs. Calcium Carbonate per ton.

*In Florida and Michigan, the heading reads "MINIMUM GUARANTEED ANALYSIS".

**In states requiring, reads "Water Soluble (K₂O)".

***In Florida, reads "not more than".

Florida Reg. No. 667-1026

USDA Reg. No. 4185-359

ACTIVE INGREDIENTS:

*Carbophenothion S-/ (p-Chlorophenylthio) methyl
0,0-diethyl phosphorodithioate

3.15%

Florida Requirement
Lbs. per Ton
63

INERT INGREDIENTS:

TOTAL

96.85%
100.00%

YELLOW

*Trithion - Reg. TM of Stauffer Chemical Co.
U.S. Pat. No. 2,793,224

**Equivalent to 8.22 Lbs. of Trithion actual per acre

FACTS ABOUT CHINCH BUGS

Chinch bugs are particularly damaging to St. Augustine grass. Grass turns yellow, then brown, withers, and dies. As bug population increases, damaged spots merge into large patches. Unless control measures are used, the entire lawn may be destroyed.

Heavy infestations may be confirmed by parting yellowed grass and making close inspection. The adult chinch bug is black with white markings in wings. If not revealed by this method, remove both ends of a large metal can, push one end 2 to 3 inches into ground where grass is yellow but not dead, and fill with water. Maintain water level above grass for 5 minutes. Chinch bugs, if present, will float to surface. If not found in first area examined, repeat test in a few additional suspect areas.

IMPORTANT. Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using this product.

CONDITIONS OF SALE. These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are

Back Panel (continued)

Bag Print
NUTRO CHINCH BUG CONTROL
Turf Food 20-5-5
Size: 15 Lbs.

GOOD LAWNS MAKE GOOD NEIGHBORS

THE BORDEN CHEMICAL COMPANY, SMITH-DOUGLASS DIVISION, NORFOLK, VA. 23501


SPREADER SETTINGS

TRUE TEMPER	4½
NUTRO AND ROTARY CYCLONE	5
AMC & MODERN	6
NUTRO & CENTRAL IMPERIAL	6
LAWN BEAUTY	7
SCOTT 35-3, 35-8	7
SEARS AND JACKSON 20-26	7


These settings are approximate. Settings vary with model, age and condition of spreader.

NUTRO CHINCH BUG CONTROL
(R)

Turf Food



Front Panel



BORDEN
CHEMICAL

NUTRO
(
CHINCH BUG CO
Turf Foo

for exclusive use

CONTROLS CHINCH BUGS &
Feeds grass back to

USDA Reg. No. 4185-359

ACTIVE INGREDIENTS:

*Carbophenothion S-l (p-Chlorophenylthio)
phosphorodithioate

INERT INGREDIENTS:

TOTAL

*Trithion - Reg. TM of Stauffer Chemical Co.
U. S. Pat. No. 2,793,224

**Equivalent to 8.22 Lbs. of Trithion actual p

NET WT.

TREATS & FEEDS
2,500 SQ. FT.
\$5.95

NUTRO CHINCH BUG CONT
(R)

YELLOW

CAUTION: KEEP OUT OF REACH OF CHILDREN AND
cautions

Back Panel

BORDEN
CHEMICAL
NUTRO
(
CHINCH BUG CO
Turf Foo

NET WT.

20 - 5 -

Bag Print
NUTRO CHINCH BUG CONTROL
Turf Food 20-5-5
Size: 15 Lbs.

Back Panel (continued)

followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. The Borden Chemical Co. and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks.

WARRANTY AND LIMITATIONS OF DAMAGES. The Borden Chemical Co. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale; subject to the risks referred to therein. The Borden Chemical Co. makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of the Borden Chemical Co. is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

DIRECTIONS FOR USE ON LAWN AREAS ONLY

YELLOW

Controls Chinch Bugs and Bermuda Mites

WHEN TO APPLY: Use immediately upon evidence that chinch bugs are at work. As a preventative, use as soon as weather is consistently above 80 degrees and repeat application at two month intervals, depending on length of chinch bug season in your area.

HOW TO APPLY: Use a fertilizer spreader and apply this bag to 2,500 sq. ft. (50' x 50') of lawn. Settings below are approximate for applying contents of this bag at suggested rate of 6 pounds per 1,000 sq. ft. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed.

Immediately after application, drench area with water or sprinkle thoroughly. Do not allow persons or pets on treated lawn until grass is completely dry.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION - HAZARDOUS IF SWALLOWED. Avoid breathing dust. Wear a dust respirator. Wear gloves. Avoid contact with skin and eyes. Wash hands with soap and water before eating, smoking, and after using. Wash all contaminated clothing with soap and hot water before re-use. Do not apply when weather conditions favor drift of dust from areas treated. Do not contaminate feed and foodstuffs.

This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be killed.

Keep children and pets off treated area until treated areas are well watered and completely dried.

STORE OUT OF REACH OF CHILDREN. Trithion, upon repeated, prolonged and careless use, may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL.

DESTROY THIS BAG BY BURNING WHEN EMPTY. Avoid exposure of persons or animals to smoke.

YELLOW

4-25-68


4185-359

Bag Print
NUTRO CHINCH BUG CONTROL
Turf Food 20-5-5
Size: 15 lbs.

Back Panel (continued)

GUARANTEED ANALYSIS*

TOTAL NITROGEN (N)	20.00%
1.0% Ammoniacal Nitrogen	
13.5% Urea Nitrogen	
5.5% Water Insoluble Nitrogen	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)	5.00%
**SOLUBLE POTASH (K ₂ O)	5.00%
Total Available Primary Plant Food - 30.00%	
***Chlorine (Cl)	1.00%
Sulphur (S)	1.00%
Iron (Fe)	0.2%



Derived from - Primary Plant Foods: Anhydrous Ammonia, Urea, Urea-
Formaldehyde, Phosphoric Acid, Muriate of Potash and Sulphate of Potash.

Derived from - Secondary Plant Foods: Sulphate of Potash and Iron
Phosphate.

Potential acidity equivalent to 800 lbs. Calcium Carbonate per ton.

*In Florida and Michigan, the heading reads "MINIMUM GUARANTEED ANALYSIS".

**In states requiring, reads "Water Soluble (K₂O)".

***In Florida, reads "not more than".

Florida Reg. No. 667-1026

USDA Reg. No. 4185-359

ACTIVE INGREDIENTS:

*Carbophenothion S-[(p-Chlorophenylthio) methyl]
0,0-diethyl phosphorodithioate

3.15%**

Florida Requirement
Lbs. per Ton
63

INERT INGREDIENTS:

TOTAL

96.85%
100.00%

YELLOW

*Trithion - Reg. TM of Stauffer Chemical Co.
U.S. Pat. No. 2,793,224

**Equivalent to 8.22 Lbs. of Trithion actual per acre

FACTS ABOUT CHINCH BUGS

Chinch bugs are particularly damaging to St. Augustine grass. Grass turns yellow, then brown, withers, and dies. As bug population increases, damaged spots merge into large patches. Unless control measures are used, the entire lawn may be destroyed.

Heavy infestations may be confirmed by parting yellowed grass and making close inspection. The adult chinch bug is black with white markings in wings. If not revealed by this method, remove both ends of a large metal can, push one end 2 to 3 inches into ground where grass is yellow but not dead, and fill with water. Maintain water level above grass for 5 minutes. Chinch bugs, if present, will float to surface. If not found in first area examined, repeat test in a few additional suspect areas.

IMPORTANT. Read these entire Directions and Conditions of Sale, including the Warranty and Limitation of Damages provision, before using this product.

CONDITIONS OF SALE. These Directions for Use reflect the opinion of experts based on tests of effectiveness, of toxicity to plants and laboratory animals and of residues on or in food and feed under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are

Back Panel (continued)

Bag Print
NUTRO CHINCH BUG CONTROL
Turf Food 20-5-5
Size: 15 Lbs.

GOOD LAWNS MAKE GOOD NEIGHBORS

THE BORDEN CHEMICAL COMPANY, SMITH-DOUGLASS DIVISION, NORFOLK, VA. 23501

SPREADER SETTINGS

TRUE TEMPER	4½
NUTRO AND ROTARY CYCLONE	5
AMC & MODERN	6
NUTRO & CENTRAL IMPERIAL	6
LAWN BEAUTY	7
SCOTT 35-3, 35-8	7
SEARS AND JACKSON 20-26	7

These settings are approximate. Settings vary with model, age and condition of spreader.

NUTRO CHINCH BUG CONTROL
(R)

Turf Food

