

This is a one panel label measuring 8" x 9". Net contents will appear on container.

[4185-363]

SMITH-DOUGLASS  
ROTENONE 1% DUST

ACTIVE INGREDIENTS:

Rotenone ..... 1.0%

Other cube resins ..... 2.0%

INERT INGREDIENTS: ..... 97.0%

Total ..... 100.0%

CAUTION

Avoid inhalation of dust. Do not dust fruits or vegetables within 24 hours of harvest.

This product is toxic to fish and wildlife. Keep Out of lakes, streams and ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment, or disposal of wastes.

DIRECTIONS FOR USE

Rotenone dust can be used in any type of hand or power duster. Apply a light film of dust to all parts of plants, especially the underside of leaves, before insects cause damage.

Rotenone dust is effective against the following garden pests: Mexican bean beetle, bean leaf beetle, asparagus beetle, flea beetle, Colorado potato beetle, striped cucumber beetle, spotted cucumber beetle, and strawberry leaf beetle on beans, asparagus, beets, egg plant, cabbage, tomatoes, peppers, potatoes, cucumbers, squash and strawberries.


Cabbage looper, imported cabbage worm on cabbage, and caterpillars on lettuce and mustard greens

Spittle bugs on strawberries.

Raspberry and gooseberry fruitworm, imported currant worm, raspberry cane borer, and blueberry maggot on raspberry, gooseberry, currant, blueberry, and related berries. Apply to berry foliage seven days after first blossoms and repeat in 10 to 20 days.

18 Pt. Type --- CAUTION: KEEP OUT OF REACH OF CHILDREN --- 12 Pt. Type

Distributed by THE BORDEN CHEMICAL COMPANY  
Smith-Douglass Division, Norfolk, Va.



USDA REG. NO. 4185-\_\_\_\_\_