This is a one panel label measuring 8" x 9". Net contents will appear on container.

MALABIHON 7. PUST

ACTIVE INCREDIENTS:

DIRECTIONS FOR USE ON BLUEBERRIES

BLUEBERRY MACCOT: Apply at the rate of 13 lbs. per acre. Thorough full-coverage applications should be made. Repeat applications as necessary.

PLUM CURCULIO & SHARP-NOSED LLAFHOPPER: Apply at the rate of 26 2/3 to 33 1/3 lbs. per acre. Apply when largest berries of earliest varieties are 1/4 inch in diameter and 7 to 10 days later.

JAPANESE BELTLE: Apply at the rate of 13 1/3-16, per acre. Apply when first berries turn blue and at 10 day intervals as intestation persists.

CAUTION.

Harmful if swallowed. Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin; such thoroughly after using. Avoid contactivation of feed and foodstuffs. Do not breathe dust. Do not get in eyes, on skin or on clothing. This product is pollonous to fish. Reep out of lakes, streams and ponds. Do not apply when weather conditions favor drift of dust from areas treated. Apply this product only as specified on this label.

Dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by burying with wastes, or by burning (Keep out of smoke).

NOTICE OF WAPPANTY: The Borden Chemical Company, Smith-Douglass Division, warrants only that the contents of the container correspond to the analysis printed on the container, lab low term BHLIPY AND THERE ARE NO WARRANTHES THAT INTEND BEYOND SUCH AVALISTS. Buyer assumes all risks of u.e, storage or handling, whether in accordance with directions or not.

14 Pt. Type --- CAUTION: KIEP OUT OF REACH OF CHILDREN --- 10 Pt. Type

See other caution on this label

Distributed by THE BORDEN CHEMICAL COMPANY Smith-Douglass Division, Nortolk, Va.

ACCEPTED WITH COMMENTS

USDA Reg. No. 4185-