

This is a one panel label measuring 8" x 9". Net contents will appear on container.

100% DDT
MALATHION 10-10-51

3. DIRECTIONS FOR USE:

Malathion 10-10-51 is a contact insecticide of the organophosphorus type. It is effective against a wide range of insects. It is also effective against mites and ticks. It is also effective against the following insects:

.....	10.00
.....	9.00
Total	19.00

DIRECTIONS FOR USE ON BLUEBERRIES

BLUEBERRY MAGGOT: Apply at the rate of 10 lbs. per acre. Thorough full-coverage applications should be made. Repeat application as necessary.

PLUM CURCULIONID FRUIT-BORING MOTHS: Apply at the rate of 20 to 25 lbs. per acre. Apply when first berries of earliest varieties are 1/4 inch in diameter and 7 to 10 days later.

JAPANESE BEETLE: Apply at the rate of 10 lbs. per acre. Apply when first berries turn blue and at 10 day intervals as infestation persists.

CAUTION

Harmful if swallowed. Avoid prolonged breathing of dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after use. Avoid contamination of food and feedstuffs. Do not breathe dust. Do not get it in eyes, on skin or on clothing. This product is poisonous to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift or dust from areas treated. Apply this product only as directed on this label.

Dispose of waste by burying in non-cropland area away from water supplies. Containers should be disposed of by burning (with care, or by burning, away from water supplies).

NOTICE OF WARRANTY: The Dorr Chemical Company, with-houlth Division, warrants only that the contents of the container correspond to the label printed on the container, and that the product is as described on the label. NO OTHER WARRANTY IS MADE. ANYONE USING THIS PRODUCT AT HIS OWN RISK AND WITHOUT LIABILITY TO THE DORR CHEMICAL COMPANY.

14 Pt. Type --- CAUTION: --- 14 Pt. Type

See other caution on this label

Distributed by THE DORR CHEMICAL COMPANY
with-houlth Division, Norfolk, Va.