

4185-438

**RASPBERRY, BLACKBERRY AND DEWBERRY** — Anthracnose, use in preblossom stage (when canes are 6 to 8 inches tall) and at first cover (right after petal fall).

**STRAWBERRY** — Botrytis fruit rot, leaf spot. Begin applications when new growth starts and repeat at 7 day intervals as necessary.

**CAUTION**

**HARMFUL IF INHALED OR SWALLOWED.** Do not take internally. Avoid prolonged or repeated breathing of fumes. Skin contact may be harmful; avoid prolonged or repeated contact with skin. Wash hands before eating. Take shower or bath after using. Wear regular long sleeved work clothing.

**KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

Avoid contamination of feed and foodstuffs. Remove any residues at harvest by washing or peeling. Dispose of container promptly after it is empty. This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.


**NOTICE:** Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but Seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty, express or implied, except as herein stated. Seller's liability is limited to the purchase price of the product and this shall be the buyers only remedy. Buyer assumes the risk of use of this product contrary to label instructions, carelessly, or under conditions not intended by or not reasonably foreseeable to Seller.

DISTRIBUTED BY  
SMITH DOUGLASS/DIVISION OF BORDEN CHEMICAL  
BORDEN INC  
COLUMBUS, OHIO 43203

**BORDEN** *Nutro* \$1.95

# Tomato Vegetable Spray

Controls tomato blight, mildew, flea beetles, hornworms and certain other insects and diseases of tomatoes and vegetables.



**ACCEPTED**  
Sept. 24, 1969  
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 4185-438 SUBJECT TO ATTACHED COMMENTS.

USDA Reg. No. 4185-438

**ACTIVE INGREDIENTS:**

|                               |                |
|-------------------------------|----------------|
| Pyrethrins                    | 0.025%         |
| Rotenone                      | 0.128%         |
| Other Cube Extractives        | 0.256%         |
| *Piperonyl Butoxide Technical | 0.256%         |
| **Captan                      | 0.120%         |
| <b>INERT INGREDIENTS</b>      | <b>99.215%</b> |

**TOTAL** 100.00%  
\*Consists of 0.265% (butyl carbonyl) (6-propyl piperonyl) ether and 0.051% related compounds.  
\*\*(N-Trichloromethyl-mercapto-4-cyclohexene-1,2-dicarboximide)

**NET WT. 1 LB. (16 OZ.)**

**CAUTION: KEEP OUT OF REACH OF CHILDREN AND PETS**

**DIRECTIONS:** Apply in early morning or late evening when temperature is low and air is still. Apply so all exposed parts are contacted, especially upper and lower leaf surfaces. Do not wet plants when applying spray. Hold aerosol can 12" away from plant. Do not treat during periods of excessively high temperatures (above 90 degrees Fahrenheit). This might cause damage to foliage. Start application at first sign of disease or insects—repeat at 7 to 10 day intervals, as necessary. Do not apply within seven days of harvest.

**INSECTS AND DISEASES CONTROLLED**

**TOMATOES**—Early blight, late blight, gray leaf spot, septoria leaf spot, Mites, tomato fruitworm, flea beetle, tomato hornworm and stinkbugs.

**CABBAGE AND CAULIFLOWER**—Flea beetles, cabbage worms, harlequin bugs, corn earworm, leaf spot.

**CUCUMBERS, CANTALOUPE, SUMMER AND WINTER SQUASH AND WATERMELON**—Angular leaf spot, anthracnose, downy mildew, Pickle worms, squash bugs, flea beetles, cucumber beetles.

**LETTUCE**—Aphids, leafhoppers, armyworms, stinkbugs.

**PEPPERS**—Mites, tomato fruitworm, flea beetles, fall armyworm, leafhoppers, lygus bug.