


CHLORDANE 10% GRANULAR



ACTIVE INGREDIENTS:	
Technical Chlordane*	10%
INERT INGREDIENTS:	90%
TOTAL	100%

Equivalent to 6% Octachloro-4, 7-methanotetrahydroindane & 4% related compounds.

4185-455 DIRECTIONS FOR USE

For Control of Soil Insects in Corn (Field and Sweet): Broadcast Application: Make broadcast application of 26 lbs. Chlordane 10% Granular per acre for control of the soil insect complex including wireworms, white grubs, seed corn beetles and white-fringed beetles. For the control of corn rootworms (Western, Southern, Northern) and cutworms, apply 26 pounds of Chlordane 10% Granular per acre. For the control of cornfield ants and cornroot aphids, use 25 lbs. Chlordane 10% Granular per acre. Apply Chlordane after field is plowed. Disc in promptly. Use the higher rate on heavier soils and the lower rate on light sandy soils. Use the higher rates for heavy infestation of cutworms.

Row Application: To control corn rootworms (Western, Southern, Northern), cornfield ants and corn root aphids, apply 10 lbs. Chlordane 10% Granular at planting time in the furrow, behind the planter shoe and ahead of the press wheel. To control wireworms and white grubs, apply 25 pounds Chlordane 10% Granular per acre in the row at planting time. Consult State Agricultural Experiment Stations for detailed information.

For Insect Control on Ornamental Turf, Nursery and Greenhouse Soils: For the control of ants, armyworms, boxelder bugs, chinch bug, crickets, European earwigs, grasshoppers, Japanese beetle grubs, mosquitoes, rose chaffer, sod webworm, spiders, ticks, white-fringed beetle, white grubs and wireworms in Turf (such as Golf Courses, Athletic Fields, Large Estate Nurseries, Cemeteries, and Park Sod) and in Nursery and Greenhouse 10% Granular per acre (4/5 lb. to each 1,000 sq. ft.) to cover surface uniformly using a lawn fertilizer spreader. Treatment should be applied in the Spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling. Do not apply to newly seeded areas.

For Insect Control on Home Lawns: For the control of the above listed insects on home lawns, apply 32 lbs. of Chlordane 10% Granular per acre (4/5 lbs. to each 1,000 sq. ft.) to cover surface uniformly using a fertilizer spreader. Treatment should be applied in the Spring, or when insects are observed to be troublesome. Immediately after treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children and pets. Do not treat when soil is wet. Apply before rain or sprinkling. Do not apply to newly seeded areas.

Potato Soil Insects: For control of Japanese beetle larvae, white fringed beetle larvae, white grubs, wireworms, apply 26 lbs. of Chlordane 10% Granular per acre. Make application broadcast prior to or at planting time and work into the top 3 or 4 inches of the soil. Increase dosage 100% on the peat or muck soils. Consult State Agricultural Experiment Stations for additional information, as the rate of application needed will vary with local conditions.

Crabgrass Control: Apply 14 lbs. Chlordane 10% Granular per 1,000 sq. ft. of lawn or turf to suppress crabgrass seedlings as germination or spouting occurs. It is for use as an early season preventive of crabgrass seed establishment. Timing of application is important and should be applied before any germination which in many areas is during the period from April 15 to May 1. Do not apply to newly seeded areas. This treatment will not burn or injure established lawns. **Spot Treatment:** For spot treatment to infested patches, apply in late Winter or early Spring prior to crabgrass seed germination using 1.4 lbs. Chlordane 10% Granular per 100 sq. ft. **Helpful Hints:** After crabgrass seedlings have been suppressed by the Chlordane treatment, control may be assisted by: 1) Mowing the lawn to 1 1/2 - 2 inches. 2) Using an adequate quantity of fertilizer to maintain active growth. After treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children or pets.

CAUTION

Harmful if swallowed. Avoid contact with skin. Avoid inhalation. Avoid contamination of feeds and foodstuffs. In case of contact with skin, wash with soap and water.

Keep children and pets off treated lawns until Chlordane has been washed into the soil by sprinkling, drenching, or rain and grass has completely dried.

ANTIDOTE: External - In case of spillage on skin, wash promptly with soap and water.

Internal - Emetic of 1 tablespoon mustard in tumbler of water. Call Physician.

This product is poisonous to fish and wildlife. Birds feeding on treated areas may be killed. Do not apply to any area not specified on the label. Keep out of lakes streams and ponds.

Dispose of wastes by burying in non-crop land away from water supplies. Containers should be disposed of by burying with wastes.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use but seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty, express or implied, except as herein stated. Seller's liability is limited to the purchase price of the product and this shall be the buyers only remedy. Buyer assumes the risk of use of this product contrary to label instructions, carelessly, or under conditions not intended by or not reasonably foreseeable to seller.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See other cautions on this label

Distributed by SMITH-DOUGLASS
Division of Borden Chemical, Borden Inc.
Norfolk, Va.

USDA Reg. No. 4185-455