

(Continued from left panel)

ANTIDOTE: If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a physician immediately if on skin. In case of contact remove contaminated clothing and immediately wash skin with soap and water.

Methyl Parathion is a cholinesterase inhibitor and treatment of severe poison should include atropine.

ATROPISE IS ANTIDOTAL

Keep in hand an emergency supply of 20 tablets grains 1 100. In case of poisonings, give 1 tablet (grs 1 100) and call physician at once. 2 PAM is also antidotal and may be administered in conjunction with atropine.

stered in conjunction with atropine. Physician: Enough atropine must be given in the shortest practical time to produce dilated pupils. Patient must be under constant supervision and additional atropine may be required without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

DISPONAL WARNING: Do not re-use the container. Completely empty the contents of the unused chemical at least 18 inches deep in an isolated location away from any flammable materials. Burn the remains in the top of the container and burn in a hot fire until all of the plastic is completely burned off the container. Extreme caution should be exercised to insure that no remains remain in the container after it has been completely empty, and that nothing flammable remains.