

GENERAL INFORMATION: Smith-Douglass Methoxychlor 2 Lb. E. C. contains 2 pounds technical methoxychlor per gallon, and is to be used as a spray with water (forming a stable emulsion) for the control of certain insects. It is suitable for high or low volume application.

DIRECTIONS

FORAGE CROPS (Alfalfa, clover, grass, soybean, cowpea and peanut forage)—Apply at the following rate using low gallonage or conventional equipment. Start applications at first sign of infestation and repeat at 7 to 14 day intervals as needed. Do not apply later than 7 days before harvest or grazing:

Leafhopper, spittlebug, 1 to 2 qts. per acre. **Alfalfa caterpillar**, flea beetles, $1\frac{1}{2}$ to 2 qts. per acre. ***Alfalfa weevil** (larvae), alfalfa webworm, fall armyworm, clover leaf weevil, soybean caterpillar, velvet bean caterpillar, 2 to 3 qts. per acre. **Cowpea curculio**, Japanese beetle, 3 qts. per acre. **Armyworm**, 3 to 4 qts. per acre.

*For pea aphid and as an alternate program for control of alfalfa weevil, and malathion (emulsifiable liquid) at the rate of 1 to $1\frac{1}{4}$ lbs. active per acre.

POTATOES—Use 2 qts. per acre for control of Colorado potato beetle, leafhoppers, and flea beetles. Repeat as necessary.


LIVESTOCK—(except dairy animals)—For control of certain external parasites.—Use as spray only. Do not use in dipping vats.

Cattle—Use 1 qt. Methoxychlor 2 Lb. E. C. to $12\frac{1}{2}$ gals. of water. Apply 2 qts. of spray per mature animal for control of hornflies; for control of short-nosed cattle lice, apply as a drenching spray.

Hogs—For control of lice, use 1 qt. Methoxychlor 2 Lb. E. C. to $12\frac{1}{2}$ gals. of water applied as a drenching spray.

SELF APPLICATOR FLY CONTROL ON CATTLE—(except dairy animals); For control of hornflies dilute 1 gal. "Methoxychlor 2 Lb. E. C." with 4 gals. of Diesel fuel or furnace type oil. Do not use any other type of oil. Use 1 gal. of solution to saturate burlap on "back rubber"; soak burlap every two to three weeks.

GRAIN STORAGE BINS—To clean up infestations in empty storage bins, use 1 gal. Methoxychlor 2 Lb. E. C. in 10 gals. of water applied at the rate of 2 gals. of spray per 1,000 sq. ft. before products are stored. This aids in control of cadelle, flat grain beetle, saw-toothed grain beetle, lesser grain borer, red flour beetle, long-headed flour beetle, rice weevil, granary weevil, confused flour beetle, foreign grain beetle and hairy fungus beetle. NOTE—Do not add grain to bin for at least 24 hours or until walls have dried out thoroughly.



ACTIVE INGREDIENTS:

*Methoxychlor Technical
Heavy Aromatic Naphtha


INERT INGREDIENTS:

TOTAL

*24% methoxychlor, technical is equivalent to
2, 2-bis (p-methoxyphenyl)-1, 1, 1-trichloroeth
and 3% other isomers and reaction products.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

Distributed by SMITH-DOL
Division of Borden Chemical, Borden
Norfolk, Va.



CHLORZOLIN

Technical	24.38%
Naphtha	69.91%
.....	5.71%
TOTAL	100.00%

lor, technical is equivalent to 21%
noxyphenyl)-1, 1, 1-trichloroethane
somers and reaction products.

KEEP OUT OF THE REACH OF CHILDREN

uted by SMITH-DOUGLASS
ion of Borden Chemical, Borden Inc
Norfolk, Va.

CONTACT AND SPACE SPRAY FOR FLIES—To reduce populations of houseflies and stable flies in farm buildings, in barnyards, in alleys, on dumps and other outdoor locations, dilute 1 pt. of Methoxychlor 2 Lb. E. C. with 2½ gals. of water and apply in space by mist blower or directly to flies by hydraulic sprayer, hand sprayer or cylindrical pressure sprayer. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying.

COMPATIBILITY—Do not use this product with any spray materials which are incompatible with oil. Do not apply within 14 days of sulfur or sulfur-product applications.

CAUTION

Avoid prolonged or repeated contact with skin. Avoid prolonged or repeated breathing of spray mist. Avoid contamination of feed and foodstuffs. Keep from children.

IMPORTANT: Users should observe the usual precautions concerning application of oil containing products to crops during high temperatures, since many crops are then more susceptible to oil injury.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area. IF EXPOSED TO PROLONGED COLD, PLACE IN WARM STORAGE (50 to 80°F.) FOR SEVERAL HOURS AND AGITATE BEFORE USING.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty, express or implied, except as herein stated. Seller's liability is limited to the purchase price of the product and this shall be the buyer's only remedy. Buyer assumes the risk of use of this product contrary to label instructions, carelessly or under conditions not intended by or not reasonably foreseeable to seller.

3-71

USDA Reg. No. 412-100

6/20 15/1971
1185-492-1