

DIRECTIONS FOR MIXING WITH LIQUID FERTILIZER

Do not mix with liquid fertilizer until time of use. Agitation should be used and in no case should the mix be left standing more than one or two hours without agitation. A simple by-pass arrangement on the liquid fertilizer tank gives adequate agitation. The liquid fertilizer-chlordane mixture should be harrowed in or mixed with the soil immediately after applying.

IMPORTANT

On field and vegetable crops, make preplant broadcast or row (where applicable) application to soil after plowing and disc or incorporate into soil immediately.
 On berry crops, make application at base of plants and incorporate into soil. Do not apply when fruit is present.
 On tree fruits, make soil application only when no fruit is present.
 Do not graze small grain grown on treated soil.
 Where a range of rates is given, use the higher rate of Chlordane 4 Lb. LF on muck or highly organic soils.

DIRECTIONS FOR USE

CORN (Field & Sweet) To control Corn Rootworm (Northern & Southern), Corn Field Ants and Corn Root Aphids, apply 1 qt. per acre (Row) or 2 qts. per acre (Broadcast). To control ~~Seed Corn Beetle Larvae~~, Seed Corn Maggot, Cutworms, White Grubs and Wireworms, apply 2 qts. per acre (Row) or 2 1/2-4 qts. per acre (Broadcast). To control Japanese Beetle Larvae, apply 1 1/2-2 1/2 gals. per acre (Broadcast). To control Asiatic Garden Beetle Larvae, apply 2 1/2-5 qts. per acre (Broadcast). To control Green June Beetle Larvae, apply 3-5 qts. per acre (Broadcast). To control White-fringed Beetle Larvae, apply 2 1/2-5 qts. per acre (Broadcast).

PEANUTS To control Seed Corn Beetle Larvae, Seed Corn Maggot, Cutworms, White Grubs and Wireworms, apply 2 qts. per acre (Row) or 2 1/2-4 qts. per acre (Broadcast). To control Japanese Beetle Larvae, apply 1 1/2-2 1/2 gal. per acre (Broadcast). To control Asiatic Garden Beetle Larvae, apply 2 1/2-5 qts. per acre (Broadcast). To control Green June Beetle Larvae, apply 3-5 qts. per acre (Broadcast). To control White-fringed Beetle Larvae, apply 2 1/2-5 qts. per acre (Broadcast).

TOBACCO Apply preplant only to control Seed Corn Beetle Larvae, Seed Corn Maggot, Cutworms, White Grubs and Wireworms, apply 4 qts. per acre (Broadcast). To control Japanese Beetle Larvae, apply 1 1/2-2 1/2 gals. per acre (Broadcast). To control Asiatic Garden Beetle Larvae, apply 4 qts. per acre (Broadcast). To control Green June Beetle Larvae, apply 4 qts. per acre (Broadcast). To control White-fringed Beetle Larvae, apply 4 qts. per acre (Broadcast).

VEGETABLE CROPS (Beans, Beets, Black-eyed Peas, Broccoli, Brussels Sprouts, Cabbage, Cantaloupe, Cauliflower, Celery, Collards, Cowpeas, Cucumbers, Eggplant, Kale, Kohlrabi, Lettuce, Melons, Okra, Onions, Peas, Peppers, Pimentos, Potatoes, Radishes, Rutabagas, Squash, Sweet Potatoes, Tomatoes, Turnips, Watermelons) To control Cutworms, Wireworms and White Grubs, apply 2 qts. per acre (Row) or 2 1/2-4 qts. per acre (Broadcast). To control Asiatic Garden Beetle Larvae and White-fringed Beetle Larvae, apply 2 1/2-5 qts. per acre (Broadcast). To control Japanese Beetle Larvae, apply 1 1/2-2 1/2 gals. per acre (Broadcast). To control Flea Beetle Larvae (potatoes and sweet potatoes), apply 2 1/2-5 qts. per acre (Broadcast).



CHLORDANE 4 LB. LF

Contains 4 lbs. Chlordane per Gallon

ACTIVE INGREDIENTS	
Technical Chlordane*	43.69
Xylene	51.11
INERT INGREDIENTS	
	5.00
Total	100.00

*Equivalent to 26.33% Octachloro-1,7-methanotetrahydroindane and 17.56% related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Antidote and Other Cautions on Side Panel

Distributed By SMITH-DOUGLASS
 Division of Borden Chemical, Borden Inc.
 Norfolk, Va. 23501



CHLORDANE 15.1

Contains 4 lbs. Chlordane per Gallon

ACTIVE INGREDIENTS	
Chlordane (90% pure)	43.80
Xylene	1.11
INERT INGREDIENTS	5.09
	<hr/> 100.00

*Equivalent to 26.33% Dieldrin, 1.0% DDT, 1.0% DDT, 1.0% DDT, and 17.56% related compounds.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See Antidote and Other Cautions on Side Panel

Distributed By SMITH-DOUGLASS
 Division of Borden Chemical, Borden Inc.
 Norfolk, Va. 23501

FRUIT CROPS: Apples, Apricot, Cherries, Grapes, Nectarines, Peaches, Pears, Plums, Prunes, Quinces, Raspberries, Strawberries. To control White Grubs, Cutworms and Wireworms apply 1-2 qts per acre (Broadcast). To control Japanese Beetle Larvae apply 2 1/2 qts per acre (Broadcast). To control Asiatic Garden Beetle Larvae apply 2 1/2-5 qts per acre (Broadcast). To control Green June Beetle Larvae apply 3-5 qts per acre (Broadcast). To control White-fringed Beetle Larvae apply 2 1/2-5 qts per acre (Broadcast).

BERRY CROPS: Blackberries, Huckleberries, Loganberries, Raspberries, Strawberries. To control Aphids apply 1-2 qts per acre (Broadcast). To control White Grubs, Cutworms and Wireworms apply 2 1/2-4 qts per acre (Broadcast). To control Japanese Beetle Larvae apply 2 1/2 qts per acre (Broadcast). To control Asiatic Garden Beetle Larvae apply 2 1/2-5 qts per acre (Broadcast). To control Green June Beetle Larvae apply 3-5 qts per acre (Broadcast). To control White-fringed Beetle Larvae apply 2 1/2-5 qts per acre (Broadcast). (Blackberries, Blueberries, Raspberries, Strawberries) To control Root Weevil Larvae apply 2 1/2 qts per acre (Broadcast). To control Rootworms, apply 1-4 2 1/2 qts per acre (Broadcast).

SMALL GRAIN: (Barley, Oats, Rye, Wheat) To control Wireworms and White Grubs apply 1 1/2 qts per acre (Broadcast). To control Japanese Beetle Larvae apply 2 1/2 qts per acre (Broadcast).

SUGAR BEETS: To control Wireworms and White Grubs apply 1 1/2 qts per acre (Broadcast).

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and farmstuffs. Keep out of reach of children.

Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be harmed. Do not apply when weather conditions favor drift from areas treated.

Do not use this product in any manner not specified on this label.

Do not use near open flame or heat.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning (keep out of smoke).

ANTIDOTE

External—In case of spillage on skin, wash promptly with soap and water.

Internal—Emetic of one tablespoonful of mustard in tumbler of water. Call a physician.

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions or in normal conditions of use but seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty express or implied except as herein stated and seller's liability is limited to the purchase price of the product and this shall be the buyer's only remedy. Buyer assumes the risk of use of this product contrary to label instructions carelessly or under conditions not intended by or not reasonably foreseeable to seller.