


4-4-73
570



ACTIVE INGREDIENT
Zineb (Zinc Ethylene Bisdiethyl Carbamate)
INERT INGREDIENTS

93.5%
93.5%
100.0%

TOTAL

DIRECTIONS

ZINEB DUST may be applied to most crops to control fungi when dusting conditions are favorable. Begin applications soon after plants are above ground or when disease conditions appear. Apply at 5 to 10 day intervals. If severe disease conditions occur, use 20 to 46 pounds of dust per acre depending on the crop size of the plants and severity of disease. Power dusters are recommended and operator should be guided by the adjustment of nozzle spacing and the spray pattern to insure adequate coverage of all parts of the plant. 5 to 10 pounds of total dust per acre is recommended for drip line applications. However, for more than 46 pounds of this material should be used per acre.

TABACCO PLANT BED: To control tobacco budworm, apply Zineb dust to the bed top. Be certain that this material is applied to the bed top and not to the sides without further mixing. Treat the bed top with Zineb dust when the plants are the size of a finger. Dusting will be most effective when plants are making extensive growth. If the tobacco budworm is not controlled by the Zineb dusts between the first and second applications, a second application should be made after the first application.

AMOUNT OF DUST PER ACRE: For most crops, use 5 to 10 pounds of dust per acre. For tobacco, use 20 to 46 pounds of dust per acre. For other crops, use 5 to 10 pounds of dust per acre.

CAUTION: Do not use this material on crops which are sensitive to Zineb dust. Do not use this material on crops which are sensitive to Zineb dust.

TOMATOES: For Early and Late Blight, Anthracnose, and Septal Leaf Spot. Do not apply within 5 days of harvest.

CUCUMBERS, MELONS, SQUASH, PUMPKINS: For Downy Mildew, Anthracnose, Angular Leaf Spot, and Leaf Blight. Do not apply within 5 days of harvest.

CELERY: For Early and Late Blight, Resin Rot, and Septal Leaf Spot. Do not apply within 5 days of harvest.

PEPPERS: For Early and Late Blight, Resin Rot, and Septal Leaf Spot.

EGGPLANTS: For control of Phytophthora, Alternaria, and Septal Leaf Spot.

CAUTION

May cause irritation to the eyes, nose, throat, and skin. Avoid contact with eyes, nose, throat, and skin.

Do not apply when weather conditions favor drift. Do not apply to crops which are sensitive to Zineb dust. Do not use this material on crops which are sensitive to Zineb dust.

NOTICE: This material is a restricted use pesticide. It may only be used by or under the direct supervision of a person who is licensed by the state to apply restricted use pesticides. This material is not to be used on crops which are sensitive to Zineb dust. Do not use this material on crops which are sensitive to Zineb dust.

CAUTION: KEEP OUT OF REACH OF CHILDREN

Warning: This product is highly toxic.
Distributed By SMITH-DOUGLASS