





DIRECTIONS FOR USE

For use on all crops except citrus, grapes, and ornamentals. Apply 1/2 pint per acre. Mix with water and spray. Do not apply to plants under stress or during periods of rapid growth. Do not apply to plants that are susceptible to frost damage. Do not apply to plants that are susceptible to leaf burn. Do not apply to plants that are susceptible to fruit drop. Do not apply to plants that are susceptible to fruit cracking. Do not apply to plants that are susceptible to fruit rot. Do not apply to plants that are susceptible to fruit decay. Do not apply to plants that are susceptible to fruit mold. Do not apply to plants that are susceptible to fruit blight. Do not apply to plants that are susceptible to fruit canker. Do not apply to plants that are susceptible to fruit scab. Do not apply to plants that are susceptible to fruit rust. Do not apply to plants that are susceptible to fruit spot. Do not apply to plants that are susceptible to fruit streak. Do not apply to plants that are susceptible to fruit blotch. Do not apply to plants that are susceptible to fruit necrosis. Do not apply to plants that are susceptible to fruit abortion. Do not apply to plants that are susceptible to fruit drop. Do not apply to plants that are susceptible to fruit cracking. Do not apply to plants that are susceptible to fruit rot. Do not apply to plants that are susceptible to fruit decay. Do not apply to plants that are susceptible to fruit mold. Do not apply to plants that are susceptible to fruit blight. Do not apply to plants that are susceptible to fruit canker. Do not apply to plants that are susceptible to fruit scab. Do not apply to plants that are susceptible to fruit rust. Do not apply to plants that are susceptible to fruit spot. Do not apply to plants that are susceptible to fruit streak. Do not apply to plants that are susceptible to fruit blotch. Do not apply to plants that are susceptible to fruit necrosis. Do not apply to plants that are susceptible to fruit abortion.

CAUTION

Do not apply to plants under stress or during periods of rapid growth. Do not apply to plants that are susceptible to frost damage. Do not apply to plants that are susceptible to leaf burn. Do not apply to plants that are susceptible to fruit drop. Do not apply to plants that are susceptible to fruit cracking. Do not apply to plants that are susceptible to fruit rot. Do not apply to plants that are susceptible to fruit decay. Do not apply to plants that are susceptible to fruit mold. Do not apply to plants that are susceptible to fruit blight. Do not apply to plants that are susceptible to fruit canker. Do not apply to plants that are susceptible to fruit scab. Do not apply to plants that are susceptible to fruit rust. Do not apply to plants that are susceptible to fruit spot. Do not apply to plants that are susceptible to fruit streak. Do not apply to plants that are susceptible to fruit blotch. Do not apply to plants that are susceptible to fruit necrosis. Do not apply to plants that are susceptible to fruit abortion.

CAUTION: KEEP OUT OF REACH OF CHILDREN
See other cautions on Side Panel

Distributed by Smith-Douglass
Division of Borden Chemical, Borden Inc.
Norfolk, Va.