

GENERAL DIRECTIONS

Prepare a spray mixture of Trump as follows:

1. Add about half the total amount of water to be used to the spray tank.
2. Start agitator or pump bypass and add proper amount of Trump.
3. Continue mixing and add remaining water.

DIRECTIONS FOR USE ON COTTON

Early Season—For control of bollweevils, fleahoppers, plant bugs and thrips, apply 1/3 gallon of Trump in 20 to 40 gallons of water per acre. Make 2 to 3 applications at 7 day intervals starting at the two-leaf stage. For aerial application or low volume ground equipment, Trump may be applied at the above rate without dilution with water.

Mid and Late Season—To control bollweevils, bollworms and cotton leafworms, make repeat applications at 5 to 7 day intervals as counts indicate they are necessary. Apply Trump at a rate of 1/2 gallon in 20 to 40 gallons of water per acre. For aerial application or low volume ground equipment, Trump may be applied at the above rate without dilution with water.

Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

DIRECTIONS FOR USE ON SOYBEANS

Apply 3 2/3 pints of Trump per acre to control army worms, bean leaf beetles, corn earworms, grasshoppers, green cloverworms, and velvetbean caterpillars. Apply suitably diluted with water. For aerial applications, apply diluted insecticide at 3 to 4 gallons per acre. Do not apply within 21 days of harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals, being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

FOR ARMYWORM CONTROL IN SMALL GRAINS

Apply 2 2/3 pints per acre to control armyworms in wheat, oats, barley or rice. Apply suitably diluted with water. With aerial application, apply in a volume of 3 to 4 gallons per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter.

WARNING

May be fatal if swallowed. Can be absorbed through skin. Do not breathe vapors or spray mist. Do not get on skin, in eyes or on clothing. Avoid contamination of feed or foodstuffs.

FIRST AID-ANTIDOTE

Internal. Give sodium chloride 15 g. (one tablespoonful of table salt) in glass of warm water and repeat until vomit is clear.

External. Wash with soap and water

**SMITH
DOUGLASS**
FARM CHEMICALS

TRUMP

ACTIVE INGREDIENTS:

Toxaphene 56.15%
Malathion (0, 0-dimethyl phosphorodithioate of
diethyl mercaptosuccinate) 14.04%
Aromatic Petroleum Derivatives (Xylene) 25.05%

INERT INGREDIENTS: 4.76%
TOTAL 100.00%

Technical Chlorinated Camphene (containing 67-69% Camphene)
(Each Gallon Contains 6 Pounds Toxaphene and 1.5 Pounds
Malathion)

WARNING: KEEP OUT OF REACH OF CHILDREN
See side panel for additional warnings

Distributed by: SMITH-DOUGLASS
Division of Borden Chemical, Borden Inc.
Norfolk, Virginia 23501

CALL A PHYSICIAN IMMEDIATELY

PHYSICIAN'S NOTE: Toxaphene produces central nervous system excitation which responds to barbituric acid derivatives. Prevent or control convulsions with barbiturates, preferably administered intravenously. If the barbiturate can be given before convulsions start, sodium pentobarbital is the drug of choice. Since toxaphene has a fairly long duration of action, a long-acting barbiturate such as phenobarbital may be used after initial control of convulsions. Refer to "Clinical Handbook on Economic Poisons", which was prepared by the U. S. Department of Health, Education, and Welfare, Public Health Service, Communicable Disease Center, Atlanta, Georgia.

ENVIRONMENTAL CAUTIONS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

USAGE CAUTIONS

Do not graze treated areas to dairy animals or animals being finished for slaughter. Not for use or storage in or around the home environment. Destroy all containers when empty by punching holes in them and burying with wastes in non-crop land away from water supplies. Do not contaminate feed or foodstuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Keep container tightly closed when not in use. Use only as directed on label.

WARRANTY

NOTICE: Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but seller makes NO WARRANTY OF MERCHANTABILITY or FITNESS FOR A PARTICULAR PURPOSE or any other warranty, express or implied, except as herein stated. Seller's liability is limited to the purchase price of the product and this shall be the buyers only remedy. Buyer assumes the risk of use of this product contrary to label instructions, carelessly, or under conditions not intended by or not reasonably foreseeable to seller.