

Robert S. Wise Co., Inc.
502 E. 33rd. North
Wichita, Kansas 67214
Phone 316-838-1451



FRONT PANEL

HOMAR

2,4-D AMINE

Contains 4 lbs. 2,4-Dichlorophenoxyacetic Acid per Gallon

ACTIVE INGREDIENTS:

Dimethylamine 2,4-Dichlorophenoxyacetate 49.6%
2,4-Dichlorophenoxyacetic Acid Equivalent 41.2%

INERT INGREDIENTS: 50.4%

(Net Contents ___ U. S. Gallons)

(12 point) CAUTION KEEP OUT OF REACH OF CHILDREN (12 point)
See other Caution Statements on Side Panel

72.32 %

Inert Ingredients by weight 50.4 water

HoMar 2,4-D Amine

LEFT PANEL

DIRECTIONS FOR USE



For best results apply 2,4-D sprays when soil and climatic conditions are conducive to rapid growth. Weeds are most susceptible when young and growing vigorously. Sprays are most effective when temperatures range from 50° to 95°F.

Mix with water by agitation in spray tank. If allowed to stand, reagitrate before use. Use at recommended rates in sufficient water for thorough weed coverage.

SMALL GRAINS (wheat, oats, barley and rye) Apply 3/4 to 1 1/2 pints per acre in the Spring from the fully tillered to the jointing stages. Late or preharvest treatment to control large weeds may be effective under certain conditions such as adequate moisture and succulent weeds. Make application of 2 to 3 pints per acre after milk stage to harvest. Do not forage or graze treated grain within 2 weeks after treatment with 2,4-D. Do not feed treated straw to livestock.

GRAIN SORGHUM (Post-emergence) Use 3/4 to 1 1/2 pints per acre when the sorghum is to 12 inches tall and secondary roots are well established. Use drop nozzles when crop is over 10 inches tall.

CORN (Pre-emergence) Use 1 to 2 quarts in 10 to 20 gallons of water per acre. Apply to soil 3 to 5 days after planting but before corn emerges. Do not use on light sandy soil. Avoid cultivation as long as possible.

(Post-emergence) Apply 1/2 to 1 pint per acre. Use lower rate for easily killed weeds and when corn is growing rapidly. Apply when weeds are up and corn is 3 or more inches tall. As soon as corn is over 10 inches use drop nozzles to direct the spray over tops of weeds but not over the corn. Rapidly growing corn is susceptible to 2,4-D injury. Do not spray corn during period after tassels begin to show until silks dry. Do not cultivate for about 2 weeks after treatment while corn is brittle.

TURF AND PASTURE WEEDS To control many broadleafed weeds use 2 to 3 pints per acre in the amount of water needed for even spraying. Apply when weeds are growing actively. Do not use on creeping grasses such as bent, except for spot spraying, nor on freshly seeded lawns or pastures until grass is well started. Most legumes are usually damaged or killed.

Dilution Equivalent For Small Equipment: One tablespoonful of this product in 1 1/2 gallons of water is equivalent to 1 quart in 100 gallons.

Manufactured by
Robert S. Wise Co., Inc.
Wichita, Kansas

USDA Reg. No. _____

5-6-69
4252-7

RIGHT PANEL

PLANTS SUSCEPTIBLE TO 2,4-D:

annual cow thistle
barnyard
buckhorn
bull thistle
burdock
carni
chicory
cirsium
cocklebur
dandelion
dock

goldenrods
ground ivy
heal-all
Jimsonweed
lamb-quarters
lawn pennywort
kochia
loose weed
mallow
morning glory
mustards
nettles

pigweed
plantains
puncture vine
ragweed
shepherds purse
sunflower
tarweed
wild radish
verch
yarrow

CAUTION: Avoid contact with skin, eyes or clothing. Harmful if swallowed. In case of contact, flush eyes with plenty of water, wash skin with soap and water. Avoid breathing spray mist. Do not reuse empty container. Destroy by burying in a safe place.

Do not apply directly to, or otherwise permit solution or spray drift to come into contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton or other desirable plants which are sensitive to 2,4-D and its formulations, and do not permit sprays containing it to drift onto them, since even minute quantities of the spray may cause injury during both growing and dormant periods. Coarse sprays are less likely to drift. Accordingly, applications by airplane and ground rigs should be carried out only when there is no hazard from drift. Do not apply by air plane in the vicinity of cotton, grapes or other desirable 2,4-D susceptible vegetation. Vapors from this product may injure susceptible plants in the vicinity.

Do not use same equipment for other purposes. If necessary to use sprayer for any other spraying, be sure to thoroughly clean all equipment with a suitable chemical cleaner.

Do not store or place near fertilizers, seeds, plants, insecticides, or fungicides. Do not contaminate irrigation ditches or water used for domestic purposes. Do not graze treated areas by dairy animals within seven days after treatment. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth.

Do not store in freezing temperatures. If allowed to freeze, remix before using. Do not use or store near heat or open flame. Be sure that the use of this product conforms to all local applicable regulations.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.