

# AMOBAM SOLUTION

\*AMOBAM is Registered Trade Mark, Fike Chemicals, Inc.

ACTIVE INGREDIENTS  
 D-aminoacetic ethylene disulfide 42%  
 INERT INGREDIENTS 58%  
 TOTAL 100%

Amobam Solution Can Be Used to Prepare Zineb When Mixed with Zinc Sulfate or Manganese Sulfate. If Manganese Sulfate is used, the mixture will contain the nutritional value of plants.

**DO NOT REUSE CONTAINER DESTROY WHEN EMPTY**

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

Do not breathe spray mist. May cause irritation of nose and throat. May cause skin irritation. Avoid contact with skin and eyes. In case of contact, flush with plenty of water. For eyes get medical attention. Harmful if swallowed. Protective clothing (long pants, long sleeve shirts, gloves, hat & boots) must be worn during mixing and loading.

### GENERAL & MIXING DIRECTIONS

Apply Amobam Solution to plants at intervals of 7 to 10 days. Do not apply within 7 days of harvest. Amobam Solution can be used to prepare Zineb when mixed with Zinc Sulfate or Manganese Sulfate. If Manganese Sulfate is used, the mixture will contain the nutritional value of plants.

**To Prepare Tank Mix Zineb**  
 1. Sift in and dissolve 1 lb. of manganese sulfate into 100 gallons of water.  
 2. Add Amobam solution 30 to 60 sec. to the tank to mix the solution.  
 3. Add Zinc Sulfate to the tank to mix the solution.

Amobam	Manganese Sulfate	Zinc Sulfate
1 gal	1 lb	1 lb
2 gal	2 lb	2 lb
3 gal	3 lb	3 lb

**FIKE CHEMICALS, INC.**

EPA Reg. No. 255-12  
 EPA Establishment 4255-WV-1

*Protective clothing (long pants, long sleeve shirts, gloves, hat & boots) must be worn during mixing and loading.*

**APPLES** Downy mildew, Cercospora, and other diseases. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**BEANS** Downy mildew and Cercospora. Start application at early blooming stage, or when the seeds are in the soil. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**BROCCOLI BRUSSEL SPROUTS, CABBAGE and CAULIFLOWER** Alternaria leaf spot and downy mildew. Apply 2 to 3 pts. at 7 to 10 day intervals. Spreader/Nozzle may be required for maximum coverage. Do not apply within 7 days of harvest.

**CARROTS** Alternaria leaf spot and Cercospora. Apply 2 to 3 pts. per acre at 7 to 10 day intervals. Do not apply within 7 days of harvest. Greater tops are to be used for food.

**CELERY** Early blight (Cercospora) and late blight (Septoria). Apply 2 to 3 pts. per acre at 7 to 10 day intervals. Remove residues by stripping, trimming and washing.

**CUCUMBERS, MELONS, SQUASH and WATERMELONS** Downy mildew, Cercospora and other diseases. Apply 2 to 3 pts. at 7 to 10 day intervals. Under severe conditions, reapplication may be necessary. For Cercospora leaf spot use Zineb only.

**GRAPES** Downy mildew. **RESTRICTED TO GREAT LAKES AREA ONLY.** Apply AMOBAM 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest. Do not apply to grapes after bloom on 7 to 10 days later. Downy mildew. Apply at the beginning and during a tank mix Zineb but continue to apply at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**LETTUCE and ENDIVE** Downy mildew. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest. Do not apply to lettuce from seed lettuce by **stripping and trimming** the lettuce heads. Do not apply to lettuce from seed lettuce by **stripping and trimming** the lettuce heads.

**PEAS** Downy mildew, Cercospora and other diseases. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**PEANUTS** Downy mildew, Cercospora and other diseases. Apply 2 to 3 pts. Amobam 2 to Sulfate incohydate at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**POTATOES** Downy mildew. Apply 2 to 3 pts. at 7 to 10 day intervals (Mush ONLY).

**SWEET CORN** Downy mildew, Cercospora and other diseases. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**SPINACH** Downy mildew, Cercospora and other diseases. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**SWEET POTATOE** Downy mildew, Cercospora and other diseases. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**TOMATOES** Downy mildew, Cercospora and other diseases. Apply 2 to 3 pts. at 7 to 10 day intervals. Do not apply within 7 days of harvest.

**ACCEPTED**  
 JAN 17 1983  
 For the Federal Insecticide, Fungicide, and Rodenticide Act  
 4255-12