


CAUTION

BAITS PREPARED WITH WARFARIN RODANTICIDES MUST BE PLACED WHERE HUMANS, LIVESTOCK, POULTRY OR PE'S WILL NOT ACCIDENTALLY EAT THEM. IN CASE BAITS ARE EATEN GIVE A TABLESPOON OF SALT IN A GLASS OF WARM WATER AND REPEAT UNTIL VOMIT FLUID IS CLEAR. CALL A PHYSICIAN IMMEDIATELY.

NOTE TO PHYSICIAN

WHEN A HUMAN HAS BEEN KNOWN TO HAVE EATEN ANY WARFARIN RODANTICIDE, BLOOD TRANSFUSIONS COMBINED WITH INTRAVENOUS INJECTIONS AND ORAL DOSES OF VITAMIN K ARE INDICATED AS IN THE CASE OF HEMORRHAGE CAUSED BY OVERDOSE OF DICUMAROL.



ROLIN®

M EXTERMINATORS

LER, WASH. 99035

NGREDIENTS

TOTAL


TOTAL

NET WILL AND COUMARIN

100.00055

EXTERMINATORS - TYLER, WASH. 99035

NET WILL 4 0Z.



GENERAL DIRECTIONS

PLACE BAITS IN LOCATIONS WHERE RATS AND MICE FEED, WATER, OR TRAVEL. BAIT PLACEMENTS ALONG WALLS, IN CORNERS AND IN CONCEALED LOCATIONS WHERE RATS AND MICE CAN FEED UNOBSERVED ARE DESIRABLE. CONTAMINATED OR SPOILED BAITS SHOULD BE REPLACED PROMPTLY. BAITING SHOULD CONTINUE UNTIL ALL SIGNS OF FEEDING HAVE STOPPED.

INSTRUCTIONS FOR USE

FOR RATS USE AS IS. PLACE BAGS IN DENS, UNDER FLOORS, RUBBISH PILES, ETC. WHERE RATS ARE KNOWN TO WORK. AN UNINTERRUPTED SUPPLY OF BAIT SHOULD BE MAINTAINED FOR AT LEAST 10 DAYS. BAIT SHOULD BE PLACED IN LOCATIONS NOT ACCESSIBLE TO CHILDREN, PETS, AND DOMESTIC ANIMALS, OR IN TAMPER-PROOF BAIT BOXES.

FOR MICE PLACE BAITS AT 8-12 FOOT INTERVALS WHERE MICE ARE KNOWN TO WORK IN TABLESPOON AMOUNTS (1/4 TO 1/2 OUNCE). AN UNINTERRUPTED SUPPLY OF BAIT SHOULD BE MAINTAINED FOR AT LEAST 15 DAYS. BAIT SHOULD BE PLACED IN LOCATIONS NOT ACCESSIBLE TO CHILDREN, PETS, AND DOMESTIC ANIMALS, OR IN TAMPER-PROOF BAIT BOXES.



CAUTION

KEEP OUT OF REACH OF CHILDREN & PETS SEE OTHER PRECAUTIONS ON BACK PANEL

ACCEPTED

FEB 19 1969

HINDER THE FEDERAL INSCOTIONS

FOR ECONOMIC POISON - NTE


ED UNDER NO.42 - SUBJEC

TO ATTACHED COMMENTS





UNFAIR TO RATS



RAT. & MOUSE KILLER