

KEEP OUT OF REACH OF
CHILDREN & PETS.

CAUTION

SEE OTHER
PRECAUTIONS ON
BACK PANEL.

REREGISTERED

ACCEPTED
JUN 2 1986
<small>U.S. Federal Insecticide, Fungicide and Rodenticide Act Registration for the pesticide Warfarin EPA Reg. No. 4421-7</small>




WATSON'S MOUSE KILLER

ACTIVE INGREDIENT:

Warfarin, 3-(alpha-Acetylbenzyl)- 4-hydroxycoumarin025%
INERT INGREDIENTS:	99.975%
TOTAL	100.000%

**READY MIXED WARFARIN
NEW ENCAPSULATED**

1 LB. NET. WT.



CAUTION

Baits prepared with Warfarin Rodenticides must be placed where humans, livestock, poultry or pets will not eat them. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN - If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD:
Keep Out of Lakes, Streams or Ponds.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS - For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS - Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, but gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE: Store in dry and secure area.

DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

R&M Exterminators, Inc.

Route 4, Box 148, Cheney, WA 99004

EPA REG. NO. 4271-7
EPA EST. 4271-WA-1

