Kills Norway Rats, Roof rats, and House Mice

Ready Mixed Warfarin Encapsulated

Active Ingredient:

[3-(Acetobenzyl)-4-hydroxyxcoumarin] 0.025%
Inert Ingredients: A.C. G. B. T. I. I.
Total 100.00%

JAN 12

KEEP OUT OF REACH OF CHILDREN the Federal Insecticide,

CAUTION

See side panel for additional BPA Reg. No. #2
precautionary statements

R&M EXTERMINATORS, INC.

24212 S. "D" STREET CHENEY, WA 99004

EPA Reg. No. 4271-7 EPA Est. No. 4271-WA-1 Water the Federal Insecticide, Fundade, and Rodenticide Act as amended, for the pesticide registered under RPA Bed No 47 7/ 77

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

FIRST AID

If swallowed:

- · Call poison control center or doctor immediately for treatment advice.
- Have a person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15–20 minutes.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Network at 1-800-858-7378 for emergency treatment information.

NOTE TO PHYSICIAN

If Ingested, administer Vitamin K1 intramuscularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean water mark.

STORAGE AND DISPOSAL

STORAGE: Store only in original container, in a dry place, inaccessible to comment and pets.

DISPOSAL: If empty: Do not reuse this container. Place in Crash or offer fo. yoling if available, if partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

DIRECTIONS FOR USE

it is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget wildlife animals to rodenticides. To help to prevent accidents:

- 1. Store product not in use in a location out of reach of children and pets.
- Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper resistant bait stations. These
 stations must be resistant to destruction by dogs and by children under six year of age, and must be used in a manner that prevents such
 children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when lifted, units must be secured
 or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially
 destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container, and unused, spoiled, and unconsumed balt as specified on this label.

ISE RESTRICTIONS: For control of Norway rats. roof rats, and house mice in and around homes, industrial buildings, and similar man-made tructures. Do not place bait in areas where there is a possibility of contaminating food or surfaces that can come in direct contact with food. Do not broadcast bait.

ELECTION OF TREATMENT AREAS: Determine areas where rats or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS; Open pack and apply 4 to 16 oz. of bait per placement, Maintain an uninterrupted supply of fresh bait for at least 10 days...

MICE: Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at 8 to 12-foot intervals. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent re-infestation, limit sources of rodent food, water, and harborage as much as possible. If re-infestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

NOTICE OF WARRANTY: Manufacturer and Seller makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements of this label.

Net Weight: 50 lbs.