

FIELD RODENT BAIT

CONTAINING
STRYCHNINE

1-1C FORMULATION



DANGER



POISON



**ACTIVE INGREDIENT:
STRYCHNINE ALKALOID**

0.5 %

INERT INGREDIENTS

99.5 %

TOTAL 100.00%

DANGER: KEEP OUT OF REACH OF CHILDREN.

DANGER: CONVULSIVE POISON. BAITS SHOULD BE PLACED IN AREAS INACCESSIBLE TO CHILDREN, PETS AND DOMESTIC ANIMALS OR IN TAMPER PROOF BAIT BOXES. POISONOUS IF SWALLOWED.

BAITS SHOULD BE PLACED IN AREAS NOT ACCESSIBLE TO CHILDREN, PETS, WILDLIFE AND DOMESTIC ANIMALS OR IN TAMPER PROOF BAIT BOXES.

ANTIDOTE: CALL A PHYSICIAN IMMEDIATELY!!

IF LESS THAN TEN MINUTES HAVE PASSED SINCE THE POISON WAS TAKEN GIVE A TABLESPOON OF SALT IN A GLASS OF WARM WATER TO REMOVE STRYCHNINE FROM THE STOMACH. TREATMENT BY THIS METHOD AFTER TEN MINUTES MAY INDUCE ONSET OF CONVULSIONS. HAVE VICTIM LIE DOWN IN A QUIET DARKENED ROOM AND KEEP HIM WARM UNTIL THE PHYSICIAN ARRIVES.

WHERE PLACEMENT OF STRYCHNINE TREATED GRAIN PRESENTS A HAZARD TO WILDLIFE, PETS OR LIVESTOCK, ALL PLACEMENTS MUST BE MADE IN OR UNDER COVER. GRAIN BAITS EXPOSED ON BARE GROUND MAY BE HAZARDOUS TO BIRDS. TREATED SEEDS EXPOSED ON SOIL SURFACES MAY BE HAZARDOUS TO BIRDS AND OTHER WILDLIFE. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES. DISPOSE OF EXCESS TREATED SEEDS OR WASTES BY BURIAL. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

BURN EMPTY BAIT CONTAINERS

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

EPA REG NO. 4271-10

NET WEIGHT 25 LBS.

FIELD RODENT BAIT

FIELD RODENT BAIT

DIRECTIONS FOR USE: (0.5% Strychnine Grain)

HOUSE MICE:

Place ten to twenty grains of bait in a small container (such as a tin can) and place it in a mouse hole. The bait should be placed in the hole so that it will be pulled into the hole. The bait should be placed in the hole so that it will be pulled into the hole.

FIELD MICE (Microtus, Pitymus)

Field Mice in orchards (Microtus, Pitymus) Part the matted grass and place a teaspoonful of bait in the surface trails or at the mouth of holes leading to underground burrows. Replace grass over runway. Make 3 to 4 bait placements under the drip line of each tree. Baiting should be conducted at the rate of 2 to 3 pounds per acre.

POCKET GOPHERS:

Remove burrow plug from the flat side of the burrow with a wire handle fork and insert a teaspoon of bait into main runway system. Care should be taken to place bait in the main runway. Burrow builders may be used for treating extensive areas. One pound will treat five to eight acres when applied by hand. One to two pounds will treat one acre when applied with burrow builder.

GROUND SQUIRRELS:

Scatter one teaspoon of bait over a square foot of clear ground within two or three feet of burrow entrance. One pound will treat ten to fifteen acres depending on animal population density.

FOR MACHINE METHOD CONTROL OF POCKET GOPHERS DIRECTIONS FOR USE

Treatments are made at 20' to 25' intervals. Between 1½ lbs. to 3 lbs. of bait per acre is required.

TREATMENT FOR FIELD MOUSE CONTROL DIRECTIONS FOR USE

Field Mice in Orchards (Microtus, Pitymus): Part the matted grass and place a teaspoonful of bait in the surface trails or at the mouth of holes leading to underground burrows. Replace grass over runway. Make 3 to 4 bait placements under the drip line of each tree. Baiting should be conducted at the rate of 2 to 3 pounds per acre.

PREPARED BY

R & M EXTERMINATORS
TYLER, WASHINGTON