

DANGER

Keep Out Of Reach Of Children.

Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

CARROLL QUAT SANITIZER

DISINFECTANT · DEODORIZER · SANITIZER

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides . . . 5.0%
n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
ethylbenzyl ammonium chlorides . . . 5.0%

INERT INGREDIENTS: 90.0%

THIS PRODUCT IS A CONCENTRATE
AND SHOULD BE DILUTED BEFORE USING.

DANGER

KEEP OUT OF THE REACH OF CHILDREN.

See other precautionary and first aid statements on left panel.

This product when used on environmental, inanimate, hard surfaces at 3-1/2 oz. per 5 gallons of water is effective against Influenza A₂, Herpes Simplex, Adenovirus type 5, and Vaccinia Viruses.

COLD DISINFECTION: For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1-1/2 ounces Sanitizer per gallon of water for 10 minutes.

EPA REG. NO. 4313-54
EPA Est. 4313-TX-1

1 U. S. GALLON
3.785 liters

ACCEPTED
23 JUL 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED UNDER 7513-54

USE DIRECTIONS

GENERAL PURPOSE: For sanitizing walls and inanimate hard surface Locker Rooms, Garbage Pails, Sink rooms, Offices, and Shower Stalls. or cloth. Sanitize with 1 ounce per water. Disinfect with 3 ounces Sanitizer Kills Staphylococcus aureus and Salmonella heavily soiled or contaminated areas recommended.

HOSPITALS: For floors, walls, and In Nursing Homes, Institutions, Pails, Telephones, and Restrooms, 1 ounces per 5 gallons of water. At this level, Sanitizer kills Pseudomonas aeruginosa. For heavily areas, a pre-cleaning step is recommended.

CARROLL

CARROLL

CARROLL

CARROLL

CARROLL

QUAT SANITIZER

DEODORIZER · SANITIZER

12, 5% C18)
ides . . . 5.0%
yl
5.0%
90.0%

This product when used on environmental, inanimate, hard surfaces at 3-1/2 oz. per 5 gallons of water is effective against Influenza A₂, Herpes Simplex, Adenovirus type 5, and Vaccinia Viruses.

COLD DISINFECTION: For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1-1/2 ounces Sanitizer per gallon of water for 10 minutes.

EPA REG. NO.4313-54
EPA Est. 4313-TX-1

1 U. S. GALLON
3.785 liters

ACCEPTED
23 JUL 1975
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED UNDER 4313-54

USE DIRECTIONS:

GENERAL PURPOSE: For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices, and Shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce Sanitizer to 4 gallons of water. Disinfect with 3 ounces Sanitizer to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

HOSPITALS: For floors, walls, and other hard surfaces. In Nursing Homes, Institutions, for Sink Tops, Garbage Pails, Telephones, and Restrooms, use Sanitizer at 3-1/2 ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level, Sanitizer will be effective against Pseudomonas aeruginosa. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

FOOD PROCESSING EQUIPMENT: For sanitization of previously cleaned food processing equipment and food utensils, dilute 1 oz. Sanitizer per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, **no potable rinse water** is required.

DAIRIES: To sanitize dairy equipment such as tanks, lines, pails, and milk cans, first clean and rinse the equipment thoroughly. Then apply solution containing 1 oz. Sanitizer to 4 gallons of water (200 ppm). At this level **no potable water rinse** is required. Follow recommendations of local Health Board.

RESTAURANT AND BAR RINSE: Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 oz. Sanitizer to 4 gallons of water. **No terminal potable water rinse required.**

CONCENTRATE
BEFORE USING.

CHILDREN.
Directions on left panel.

CARROLL

CARROLL

CARROLL

CARROLL COMPANY
2900 W. Kingsley Rd.
Garland, Texas 75041
214 278-1304

CARROLL

CARROLL