

FEB 20 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED-UNDER NO. 4330-1

**FOR PROFESSIONAL USE
CONCENTRATE — MUST BE DILUTED**

Waverly[®]
ANTI-RUST
SURGISEPT
SANITIZING SOLUTION

ACTIVE INGREDIENTS: isopropyl alcohol — 24.5%
n-alkyl (C₁₄ 50%, C₁₂ 40%, C₁₆ 10%) — 3 %
dimethyl benzyl ammonium chloride — 1.5%
tetrapotassium salt of ethylenediamine
tetraacetic acid

INERT INGREDIENTS : — 71 %
Contains Sodium Nitrite

DANGER

KEEP OUT OF REACH OF CHILDREN
Corrosive. Causes eye damage and skin irritation.
Do not get in eyes, on skin or on clothing. Pro-
tect eyes and skin when handling. Harmful if
swallowed. Avoid contamination of food.
Rinse empty container thoroughly with water and
discard it.

**SEE BACK PANEL FOR FIRST AID
AND NOTE TO PHYSICIAN INFORMATION.**

CONTENTS ONE GALLON

EPA REGISTRATION NO. 4330-1

WAVERLY BEAUTY PRODUCTS INC.
COLLEGE POINT, N.Y. 11356

**FOR PROFESSIONAL USE
CONCENTRATE — MUST BE DILUTED**

DISINFECTION DIRECTIONS

FOR BARBER AND BEAUTY SHOP EQUIPMENT:

DILUTION: 1 oz. (2 tablespoons) to 1 quart water or 4 oz.
to 1 gallon water.

USE: Completely immerse cleaned implements (free from
soap) in SURGISEPT solution for 15 seconds (or as re-
quired by local authorities — usual requirements, 10
minutes).

FOR MANICURING:

DILUTION: 1/4 oz. (1 teaspoon in 1 cup) (8 oz.) water.

USE: Soak fingers (free from soap) as usual, acts as an ex-
cellent cuticle softener.

**FOR COLD DISINFECTION OF
SURGICAL EQUIPMENT:**

DILUTION: 1 oz. (2 tablespoons) to 1 quart water or 4 oz.
to 1 gallon water.

USE: Completely immerse cleaned implements (free from
soap) in SURGISEPT solution for 10 minutes.

**DO NOT USE SOAP OR DETERGENTS
WITH SURGISEPT SOLUTION.**

**CHANGE SOLUTION ABOUT EVERY THREE DAYS (OR
MORE FREQUENTLY IF THE SOLUTION
BECOMES CLOUDY).**

DOES NOT DULL SHARP CUTTING EDGES

FIRST AID:

In case of contact, immediately flush eyes or skin with
plenty of water for at least 15 minutes. For eyes, call a
physician. Remove and wash contaminated clothing before
reuse. If swallowed, drink promptly a large quantity of milk,
egg whites, gelatin solution; or if these are not available,
drink large quantities of water. Avoid alcohol. Call a
physician.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of
gastric lavage. Measures against circulatory shock, respir-
atory depression and convulsion may be needed.