


PM 31 EPA# 4330-1 pg 202
CLF 12/5/89



FOR PROFESSIONAL USE
CONCENTRATE—MUST BE DILUTED

ONE-DER ANTISEPTIC

Sanitizing Solution with Anti-Rust for Manicures and Pedicures

and
Equipment

ACTIVE INGREDIENTS:
n-alkyl (C14 50%, C40%, C15 10%)
dimethyl benzyl ammonium chloride 0.3%

INERT INGREDIENTS: 97%

Contains Sodium Nitrite

DANGER

KEEP OUT OF REACH OF CHILDREN

Concentrated corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

Rinse empty container thoroughly with water and discard it.

SEE BACK PANEL FOR FIRST AID
AND NOTE TO PHYSICIAN INFORMATION.
CONTENTS 1/2 GALLON (1.89 Liters)



EPA EST. NO. 4330-NY-01
EPA REGISTRATION NO. 4330-01

DIRECTIONS

FOR MANICURES AND PEDICURES

DILUTION: 1/4 oz. ONE-DER (1 teaspoon) in 2 cups (5 oz.) water or 4 oz. ONE-DER to 1 gal. water.

FOR MANICURES: Soak fingers (free from soap) as usual.
Acts as an excellent cuticle softener.
Rinse.

FOR PEDICURES: Submerge and soak feet for approximately 10 minutes.
Rinse.

FOR COLD DISINFECTION OF SURGICAL EQUIPMENT AND BARBER SHOP AND BEAUTY SHOP EQUIPMENT

DILUTION: 1 oz. ONE-DER (2 tablespoons) to 1 quart water or
4 oz. ONE-DER to 1 gallon water.

USE: Completely immerse cleaned implements (free from soap) in ONE-DER solution for 15 seconds (or as required by local authorities - usual requirements 10 minutes)

DO NOT USE SOAP OR DETERGENTS
WITH ONE-DER SOLUTION

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsions may be needed.



Holtsville
New York, 11742

Not Reviewed. Registrant claims to be in accordance with
Registration of Pesticide Product Guidance for Phase 2
Exemption, Page 2.12, Active to Inert Change in Status.

of label with changes

pg. 2072

**FOR PROFESSIONAL USE
CONCENTRATE — MUST BE DILUTED**

Waverly
ANTI-RUST
SURGISEPT
SANITIZING SOLUTION

ACTIVE INGREDIENTS:
n-alkyl (C₁₄ 80%, C₁₀ 10%)
dimethyl benzyl ammonium chloride 0.3%

INERT INGREDIENTS: 97%
Contains Sodium Nitrite

DANGER

KEEP OUT OF REACH OF CHILDREN
Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.
Rinse empty container thoroughly with water and discard it.

SEE BACK PANEL FOR FIRST AID
AND NOTE TO PHYSICIAN INFORMATION.
CONTENTS 16 FL.OZ. (1 PINT)
EPA EST. NO. 4330-NY-01
EPA REGISTRATION NO. 4330-1
WAVERLY BEAUTY PRODUCTS, INC.
Hollisville, N.Y. 11742

**FOR PROFESSIONAL USE
CONCENTRATE — MUST BE DILUTED**

DISINFECTION DIRECTIONS

FOR BARRIER AND BEAUTY SHOP EQUIPMENT:
DILUTION: 1 oz. (2 tablespoons) to 1 quart water or 4 oz. to 1 gallon water

USE: Completely immerse cleaned implements (free from soap) in SURGISEPT solution for 15 seconds (or as required by local authorities — usual requirements, 10 minutes).

**FOR MANICURING, PEDICURING,
AND FOOTBATHS:**

DILUTION: 1/4 oz. (1 teaspoon) in 1 cup (8 oz.) water.
USE: Soak fingers (free from soap) as usual, acts as an excellent cuticle softener. (For footbaths soak 10 minutes.)

**FOR COLD DISINFECTION OF
SURGICAL EQUIPMENT:**

DILUTION: 1 oz. (2 tablespoons) to 1 quart water or 4 oz. to 1 gallon water.
USE: Completely immerse cleaned implements (free from soap) in SURGISEPT solution for 10 minutes.

**DO NOT USE SOAP OR DETERGENTS
WITH SURGISEPT SOLUTION.**

**CHANGE SOLUTION ABOUT EVERY THREE DAYS
(OR MORE FREQUENTLY IF THE SOLUTION
BECOMES CLOUDY.)**

DOES NOT DULL SHARP CUTTING EDGES

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Not Reviewed. Registrant claims to be in accordance with registration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.