

**FOR PROFESSIONAL USE
CONCENTRATE — MUST BE DILUTED**

**Waverly
ANTI-RUST
SURGISEPT
SANITIZING SOLUTION**

ACTIVE INGREDIENTS:

alkyl (C11 50%, C40%, C1510%)
dimethyl benzyl ammonium chloride3%
INERT INGREDIENTS97%

Contains Sodium Nitrite

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

Rinse empty container thoroughly with water and discard it.


**SEE BACK PANEL FOR FIRST AID
AND NOTE TO PHYSICIAN INFORMATION.**

CONTENTS 128 FL. OZ. (1 GALLON)

F.A. EST. NO 4330-NY-01

EPA REGISTRATION NO. 4330-1

**WAVERLY BEAUTY PRODUCTS INC.
Holtsville, NY 11742**



**FOR PROFESSIONAL USE
CONCENTRATE — MUST BE DILUTED**

DISINFECTION DIRECTIONS

FOR BARBER AND BEAUTY SHOP EQUIPMENT:

DILUTION: 1 oz. (2 tablespoons) to 1 quart water or 4 oz. to 1 gallon water.
USE: Completely immerse cleaned implements (free from soap) in SURGISEPT solution for 15 seconds (or as required by local authorities — usual requirements, 10 minutes).

FOR MANICURING, PEDICURING, AND FOOTBATHS:

DILUTION: ¼ oz. (1 teaspoon) in 1 cup (8 oz.) water.
USE: Soak fingers (free from soap) as usual, acts as an excellent cuticle softener. (For footbaths soak 10 minutes.)

FOR COLD DISINFECTION OF SURGICAL EQUIPMENT:

DILUTION: 1 oz. (2 tablespoons) to 1 quart water or 4 oz. to 1 gallon water.
USE: Completely immerse cleaned implements (free from soap) in SURGISEPT solution for 10 minutes.

**DO NOT USE SOAP OR DETERGENTS WITH SURGISEPT SOLUTION.
CHANGE SOLUTION ABOUT EVERY THREE DAYS (OR MORE FREQUENTLY IF THE
SOLUTION BECOMES CLOUDY.)
DOES NOT DULL SHARP CUTTING EDGES**

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

...lawed. For... with