

7389-3

Food processing plants, bakeries, Canneries, Meat packing plants, Sea Food establishments.

(for removing & preventing mold, for elimination of odors, for sanitizing equipment)

WALLS, FLOORS & CEILINGS:

1. Clean first with a good cleaning agent (Not soap). Rinse.
2. Spray or wipe with SANICIDE, sol'n. No. 3.
3. Do not rinse with water (floors, walls, and ceilings only).

FOOD PROCESSING EQUIPMENT:

1. Clean first with a good cleaning agent (not soap). Rinse.
2. Spray or wipe with SANICIDE sol'n. No. 1.
3. Rinse with clear water before equipment comes in- to contact with food.

DAIRIES & MILK FARMS

1. Clean utensils & equip- ment thoroughly with a good dairy cleaner (not soap).
2. Rinse with clear water.
3. Sanitize with SANICIDE solution No. 1.
4. Rinse with clear water before utensils come into contact with milk. SANICIDE must not be used as an antiseptic in milk.

Quality products C O - J Y Janitor supplies

SANICIDE

Sanitizer & Deodorizer

USDA Reg. No. 4889-14

Active Ingredients: Alkyl (C₈-8%, C₁₀-7%, C₁₂-48%, C₁₄-17%, C₁₆-9%, C₁₈-11%) benzylidimethyl ammonium chloride - - - - - 10%
 Tetrasodium ethylenediamine tetraacetate - - - - - 1%
 Inert Ingredients: - - - - - 89%

Phenol Coefficient: A.S.T.C. 20° C.

50 vs. Salmonella typhosa
80 vs. Staphylococcus aureus

CAUTION: KEEP OUT OF THE REACH OF CHILDREN. Harmful if swallowed. Avoid contamination of food. Avoid contact with skin or eyes. In case of contact flush with cool water. If irritation persists see a physician.

Do not use in conjunction with soap or synthetic detergents. Materials to be disinfected should be first washed and rinsed thoroughly.

Exclusive distributor: **SANICIDE SUPPLY COMPANY**
Seattle, Washington

Net contents 5 gal.

Restuarants, Fountains, Bars & Taverns

1. Clean utensils with good cleaning agent (not soap).
2. Rinse in clear water.
3. Sanitize by a 2 min. dip in SANICIDE, sol'n. No. 1. Rinse.

Hospitals, Veterinaries & Household Sanitizing.

1. Clean first with a good cleaning agent (not soap). Rinse.
2. Sanitize with SANICIDE solution No. 3.

DIRECTIONS FOR PREPARING SANITIZING SOLUTIONS

Solution No. 1 (200 ppm active) 1/500 Stir 1 oz SANICIDE in 4 gal. water.
 Solution No. 2 (400 ppm active) 1/250 Stir 1 oz SANICIDE in 2 gal. water.
 Solution No. 3 (800 ppm active) 1/125 Stir 1 oz SANICIDE in 1 gal. water.
 Two tablespoons = 1 oz.

Test: Use Stone-Marshall test papers as directed to test strength of sanitizing sol'n in use.

Non-inflammable, non-volatile, non-irritating, colorless, non-poisonous when in use dilutions. Contains no phenol, iodine, chlorine, mercury or other metals.