

Quaternary Type

MICROBICIDIAL PROPERTIES IN A USE-DILUTION CONFIRMATION TEST		
Organism	Dilution	Active Germicide
Staphylococcus aureus	1:200	575 ppm
Salmonella typhimurialis	1:200	575 ppm
Trichophyton interdigitales	1:200	575 ppm
Pseudomonas aeruginosa	1:200	575 ppm

Application	Use-Solution		Dilution	Active Germicide
	Per Gallon	Per 5 Gallons		
General disinfection of walls, lightly soiled floors	2/3 oz	3 oz.	1:200	575 ppm
General disinfection of heavily used floors, rooms and restrooms	1 1/3 oz.	7 oz.	1:100	1150 ppm
High contamination areas, heavily soiled floors & rooms	3 oz.	15 oz.	1:50	2300 ppm

DEOTERG L

GERMICIDAL DETERGENT


A wide spectrum germicide, fungicide and deodorant, combined with an efficient detergent... developed for use in hospitals. Recommended for use in general cleaning, disinfecting and deodorizing of hospital areas. Phenol Coefficient, AOAC Method: Staphylococcus aureus 40; Salmonella typhosa 73

ACTIVE INGREDIENTS:

- Didecyl Dimethyl ammonium chloride. 6.00%
- Alkyl (C₁₄-50%, C₁₂-40%, C₁₆-10%) Dimethyl Benzylammonium chloride. 5.50%
- Tetrasodium ethylene-diaminetetraacetate 4.00%
- Sodium Carbonate. 2.00%
- Essential Oils. 0.75%
- INERT INGREDIENTS: 81.75%**

DANGER: Keep Out Of The Reach Of Children Causes severe eye damage on skin, or on clothing. Wear goggles and rubber gloves when handling. Avoid ingestion of food. Rinse out empty container and discard it.
FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water. If swallowed, drink large amounts of water. If necessary, consult a physician immediately. Note to physician: Probable mucosal damage may contraindicate the use of certain drugs. Treatment against circulatory shock, respiratory depression and convulsion may be necessary.

ONE GALLON NET



FEDERAL INSECTICIDE
 AND FUNGICIDE ACT
 REGISTERED
 7387-66

PACE NATIONAL Corporation

500 7th Ave. South, Kirkland, Wash. 98033 • (206) 827-8711

APPLIED

Quaternary Type

MICROBICIDAL PROPERTIES		
USE-DILUTION CONFIRMATION TEST		
Organism	Dilution	Active Germicide
Staphylococcus aureus	1:200	575 ppm
Salmonella choleraesuis	1:200	575 ppm
Campylobacter interdigitales	1:200	575 ppm
Pseudomonas aeruginosa	1:200	575 ppm

Application	Use-Solution		Dilution	Active Germicide
	Per Gallon	Per 5 Gallons		
Disinfection, lightly soiled	2/3 oz.	3 oz.	1:200	575 ppm
Disinfection, heavily soiled and linens	1 1/3 oz.	7 oz.	1:100	1150 ppm
Sanitization, heavily soiled floors & rooms	3 oz.	15 oz.	1:50	2300 ppm

DEOTERG L

GERMICIDAL DETERGENT

A wide spectrum germicide, fungicide and deodorant, combined with an efficient detergent ... developed for use in hospitals. Recommended for use in general cleaning, disinfecting and deodorizing of hospital areas. Phenol Coefficient, AOAC Method: Staphylococcus aureus 80; Salmonella typhosa 73

ACTIVE INGREDIENTS:

Didecyl Dimethyl ammonium chloride. 6.00%
 Alkyl (D₁₄-50%, C₁₂-40%, C₁₆-10%) Dimethyl Benzylammonium chloride. 5.50%
 Tetrasodium ethylene-diaminetetraacetate 4.00%
 Sodium Carbonate. 2.00%
 Essential Oils. 0.75%
INERT INGREDIENTS: 81.75%

EPA REG. NO. 4389-60
 EPA EST. NO. 4389-WA-1

DIRECTIONS

1. Remove gross filth & heavy soil from surfaces prior to application of recommended use-solution.
2. Apply germicidal-detergent solution, per dilution table.
3. Scrub by mop, sponge, cloth or mechanical scrubber according to surface on which used.
4. Remove solution and entrained soil with wet vacuum pickup, mop, sponge or cloth, as best suited.

DANGER: Keep Out Of The Reach Of Children Causes severe eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Rinse out empty container and discard it.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, give promptly a large quantity of milk, egg whites, or gelatin solution; or if these are not available, drink large amounts of water. Avoid alcohol. Call a physician immediately. Note to physician: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

PRODUCT OF USA

E NATIONAL Corporation

10000 Ave. South, Kirkland, Wash. 98033 • (206) 827-8711

APPLIED SCIENCE

