

# PACIFIC CHEMICAL

500 7th Ave. South, Kirkland, Washington 98033 • (206) 827-8711

division Pace National Corp.



APPLIED SCIENCE

ACCEPTED

01 OCT 1979

FOOD, DRUG, COSMETIC, AND PESTICIDE ACT  
FEDERAL REGISTER

## MICROCIDE L-60

ALGAE, BACTERIA & FUNGI CONTROL AGENT

EPA REG. NO. 4389-75

EPA EST. NO. 4389-WA-1

### ACTIVE INGREDIENTS:

Disodium cyanodithioimidocarbamate	3.63%
Potassium N-methyldithiocarbamate	5.07%

INERT INGREDIENTS . . . . . 91.25%

TOTAL . . . . . 100.00%

This product contains 0.77 lb. of active ingredients per gallon and weighs 8.81 lb. per gal.

### WARNING

**KEEP OUT OF REACH OF CHILDREN**

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**

**WARNING:** Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

**FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, give patient doses of powdered charcoal or mustard in warm water. Call a physician immediately.

**ENVIRONMENTAL HAZARDS:** This pesticide is toxic to fish. Do not discharge into lakes, streams, ponds, or public waters unless in accordance with an NPDES permit. For guidance, contact your Regional Office of the EPA.

### DIRECTIONS FOR USE

#### GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MICROCIDE L-60 is used to control algae, bacteria, and fungi in recirculating commercial and industrial water cooling towers. Prior to its use, systems must be cleaned to remove algal growth, microbiological slime, and other deposits. An initial slug addition of 6.9 to 13.9 fluid ounces of MICROCIDE L-60 per 1000 gallons of water is recommended. Repeat initial dosage until control is evident.

Subsequent slug additions of 2.3 to 13.9 fluid ounces of MICROCIDE L-60 per 1000 gallons of water should be employed every 1 to 5 days, or as needed. The frequency of addition depends upon the relative amount of bleedoff and the severity of the microbiological problem. Slug additions should be made in the sump of water cooling towers.

### STORAGE AND DISPOSAL

#### PROHIBITIONS:

Do not use this product in or around residential areas, schools, hospitals, or other public buildings. Do not use this product in or around water bodies.

#### PESTICIDE DISPOSAL:

Do not use this product in or around residential areas, schools, hospitals, or other public buildings. Do not use this product in or around water bodies. Do not use this product in or around water bodies.

#### CONTAINER DISPOSAL:

Triple rinse (or equivalent) and then recycle, reuse, or dispose of in accordance with applicable regulations. Do not use this product in or around water bodies.

#### GENERAL:

Do not use this product in or around residential areas, schools, hospitals, or other public buildings. Do not use this product in or around water bodies.

55 GALLONS NET (208 LITERS)

PRODUCT OF USA