

11-16-71  
4462-25

IG-7

## DIRECTIONS

### SANITIZING DAIRY FARM PIPE LINES:

Where a separate sanitizing agent is required just before milking, sanitize line by circulating a solution of this product (1 ounce to 5 gallons of water) for 5 minutes; drain thoroughly and start milking.

### MILK PLANT SANITIZATION:


After cleaning and just prior to use, all equipment surfaces should be sanitized with a solution of this product (1 ounce to 3 gallons of water) for not less than 2 minutes. Do not rinse equipment with water after treatment.

### SWIMMING POOL CHLORINATION:

Add 1 ounce of this product to each 1000 gallons pool water as needed to maintain 0.6 to 1.0 ppm residual chlorine as determined by chlorine test kit. One cup holds 8 fluid ounces of product. The pH of the pool water should be maintained between 7.2 and 7.6.

FOR GENERAL USE IN  
DAIRY FARMS - FOOD PROCESSING PLANTS

B-301



# BEVER

**ACTIVE INGREDIENTS**

SODIUM HYPOCHLORITE 8.2%

INERT INGREDIENTS 91.8%

**B★CHLOR**A LIQUID CHLORINE  
DISINFECTANT AND GERMICIDE

KEEP OUT OF REACH OF CHILDREN

de Perio for Additional Features

**TENTS ONE GALLON**

08971

N12-13

**CAUTION: KEEP OUT OF REACH OF CHILDREN.****HARMFUL IF SWALLOWED.****SKIN AND EYE CONTACT MAY CAUSE SEVERE IRRITATION. AVOID CONTACT WITH SKIN AND EYES.**

Keep in cool, dark place, as light and heat will reduce strength of product.

Rinse empty container thoroughly with water and discard it.

Add material only to water. May cause fire or explosion if mixed with other chemicals.

Degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine.

Manufactured By  
**Bever Chemical Co., Inc.**  
STOCKTON, CALIFORNIA 95201