

BEAVER

PRODUCTS

CAUTION:

Contains Sodium Dichloro-s-Triazine Trione
Harmful if Swallowed

If Swallowed, give 2 glasses of milk

Avoid Contact With Skin and Eyes

For Eyes: Rinse with water for at least 15 minutes,
and get medical attention.

CALL A PHYSICIAN

Contact with heat, moisture, foreign material or
contamination from any source may result in fire.
Avoid breathing dust. Keep cover closed when not
in use. Store in cool dry place.

Rinse empty container thoroughly with water and
discard it.

EPA Reg. No. 4482-39-AA

Manufactured by
BEAVER CHEMICAL COMPANY, INC.
STOCKTON, CALIFORNIA 95201

CONCENTRATE SANI-BAG

POWDERED BACTERICIDE

ACTIVE INGREDIENT:

Sodium Dichloro-s-triazinetrione..... 20%

INERT INGREDIENTS..... 80%

For Sanitizing Equipment, Utensils & Glassware

CAUTION KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions

NET CONTENTS

DIREC

Dairy Farms and

1. Rinse utensils and
of milk residue.
2. Then wash utensils
Detergent, rinse w
3. To sanitize utensils
sanitizing solution
ing ½ ounce to 5
DO NOT SOAK O

Poultry Operatio

1. Remove all poultry
crates.
2. Remove all litter
surfaces of facility
3. Empty all troughs,
appliances.
4. Saturate surfaces
by mixing 3 oz.
5. Ventilate buildings
not house poultry
has been absorbed
6. All treated feed re
ers, fountains, a
with potable water

Institutions — Fo

1. Wash all articles t
2. Rinse thoroughly, t
solution of this pr
sanitizer to 5 gallo
not less than 30 se

BEAVER

PRODUCTS

CONCENTRATE SANI-BAG

POWDERED BACTERICIDE

ACTIVE INGREDIENT:

Sodium Di-chloro-s-triazinetrione.....20%

INERT INGREDIENTS.....80%

For Sanitizing Equipment, Utensils & Glassware

CAUTION KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions

NET CONTENTS

DIRECTIONS FOR USE

Dairy Farms and Plants

1. Rinse utensils and equipment with clear water until free of milk residue.
2. Then wash utensils and equipment with a recommended Detergent, rinse with clear water, then sanitize.
3. To sanitize utensils and equipment before use, make a sanitizing solution (100 ppm of available chlorine) by mixing ½ ounce to 5 gallons of cold or warm (Not Hot) water. DO NOT SOAK OVER NIGHT.

Poultry Operations

1. Remove all poultry and feeds from premises, trucks, coops, crates.
2. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Saturate surfaces with (600 P.P.M. available chlorine) by mixing 3 oz. to 5 gals. water.
5. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried.
6. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be thoroughly rinsed with potable water prior to reuse.

Institutions — Food Industry

1. Wash all articles to be sanitized with suitable Detergent.
2. Rinse thoroughly, then sanitize articles by immersing in a solution of this product (100 ppm), mixing ½ ounce of sanitizer to 5 gallons of cold or warm water (Not Hot) for not less than 30 seconds. Air dry.

53
ACCEPTED

SEP 6 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CIDE ACT
FOR EXAMINING PURPOSES
ED UNDER NO. 142-39 SUBJECT
TO ATTACHED COMMENTS.

CAUTION:

Contains Sodium Dichloro-s-Triazine Trione

Harmful If Swallowed

If Swallowed, give 2 glasses of milk

Avoid Contact With Skin and Eyes

**For Eyes: Rinse with water for at least 15 minutes,
and get medical attention.**

CALL A PHYSICIAN

Contact with heat, moisture, foreign material or contamination from any source may result in fire. Avoid breathing dust. Keep cover closed when not in use. Store in cool dry place.

Rinse empty container thoroughly with water and discard it.

EPA Reg. No. 4462-39-AA

Manufactured by
BEAVER CHEMICAL COMPANY, INC.
STOCKTON, CALIFORNIA 95201

**For S
CA**

PRODUCTS

CONCENTRATE SANI-BAC

POWDERED BACTERICIDE

ACTIVE INGREDIENT:

Sodium Dichloro-s-triazinetrione..... 20%

INERT INGREDIENTS..... 80%

For Sanitizing Equipment, Utensils & Glassware

CAUTION KEEP OUT OF REACH OF CHILDREN

See Side Panel for Additional Cautions

NET CONTENTS

of milk residue.

2. Then wash utensils and equipment with a recommended Detergent, rinse with clear water, then sanitize.
3. To sanitize utensils and equipment before use, make a sanitizing solution (100 ppm of available chlorine) by mixing ½ ounce to 5 gallons of cold or warm (Not Hot) water. DO NOT SOAK OVER NIGHT.

Poultry Operations

1. Remove all poultry and feeds from premises, trucks, coops, crates.
2. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Saturate surfaces with (600 P.P.M. available chlorine) by mixing 3 oz. to 5 gals. water.
5. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried.
6. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be thoroughly rinsed with potable water prior to reuse.

Institutions — Food Industry

1. Wash all articles to be sanitized with suitable Detergent.
2. Rinse thoroughly, then sanitize articles by immersing in a solution of this product (100 ppm), mixing ½ ounce of sanitizer to 5 gallons of cold or warm water (Not Hot) for not less than 30 seconds. Air dry.

