

USE DIRECTIONS

FOR CLEANING

Lightly soiled areas - surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of BEAVER SAN per gallon of water. Rinse cleaned area with fresh water.

Heavily soiled areas - mop or brush soiled surface with a solution containing 2 - 3 ounces of BEAVER SAN per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, mop or brush surface to be treated with a solution containing 1 ounce of BEAVER-SAN per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of BEAVER-SAN per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop area with a solution containing 2 ounces in 3 gallons of water. This solution will destroy *Trichophyton interdigitale*, a causative agent of athlete's foot.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 ounces of BEAVER-SAN per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing 1 ounce of BEAVER SAN per gallon of water. If

BEAVER SAN

Phenol Coeff: S. aureus - 38
S. typhosa - 42

DISINFECTANT-CLEANER-SANITIZER-FUNGICIDE-DEODORANT

ACTIVE INGREDIENTS

n-Alkyl (50% C₁₄, 40% C₁₂, 10% C₁₆) dimethyl benzyl ammonium chloride 5.0%

Tetrasodium salt of ethylene diamine tetracetic acid 2.3%

Sodium carbonate 2.0%

INERT INGREDIENTS 90.7%

EPA Reg. No. 4462 - 55

DANGER

KEEP OUT OF REACH OF CHILDREN.


See antiseptic statement and additional precautions on side panel.

Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

**MANUFACTURED BY
BEAVER CHEMICAL CO., INC.
1448 SHAW ROAD
STOCKTON, CAL. 95201**

NET CONTENTS

1 - GALLON



4462-55

surface is heavily soiled, use BEAVER-SAN at a rate of 2 - 3 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 1/2 ounce of BEAVER-SAN per gallon of water. No rinse required after sanitization.

DEODORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of BEAVER-SAN per gallon of water. BEAVER-SAN is an effective antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, BEAVER-SAN will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial-antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

See left panel for use directions.

DANGER

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.