



SCREW WORM SPRAY

Wound Protectant - Protects Wounds from
Screw Worms and Blowfly Maggots and
Kills Screw Worms and Blowfly Maggots

For livestock of all ages, including newborn animals

ACTIVE INGREDIENTS:

Ronnal O,O-dimethyl O-(2,4,5-tri-chlorophenyl) phosphorothioate	-----	2.5%
Xylene	-----	0.5%
INERT INGREDIENTS	-----	97.0%

FOR EXTERNAL USE ONLY
LIVESTOCK REMEDY - NOT FOR HUMAN USE

WARNING: Contents Under Pressure.

CAUTION: Keep Out of Reach of Children.
Harmful if Swallowed. Injurious to Eyes. See back panel for additional Cautions.

NET WEIGHT 10 OUNCES

MORTON PHARMACEUTICALS, INC.
ETHICAL DISTRIBUTORS MEMPHIS, TENN 38108

DIRECTIONS FOR USE:
For Use on Cattle, Sheep, and Horses

TO OPERATE:

Hold outlet about 3 inches from wound or infested area. Be sure opening is directed toward area to be treated. Press valve down to spray, wetting wound and about 1 inch of surrounding area thoroughly.

SCREW WORM AND BLOWFLY INFESTATIONS:

Use particular care to spray into deep pockets caused by screw worms. Repeat applications at 5 to 7 day intervals until wound is healed.

TO PREVENT SCREW WORM AND BLOWFLY INFESTATIONS:

Apply directly to all open or fresh wounds resulting from de-horning, castration, shear cuts, etc. Also treat navels of newborn animals. Freely bleeding wounds may require several applications. Repeat applications at 5 to 7 day intervals until wound is healed.

FLEECE WORM INFESTATIONS:

Spray infested areas liberally, working toward the center of infestation. Shearing or removing wool from the infested area is not necessary. Re-treat in 5 to 7 days as needed.

WARNING:

Do not puncture container. Store at room temperature (not over 120° F.) away from radiators, stoves, hot water or other heat. Never throw container into fire or incinerator when empty.

CAUTION:

Avoid contact with skin, eyes and clothing. In case of contact with human eyes, flush eyes immediately with plenty of water and get prompt medical attention. Avoid breathing spray mists. Wash hands thoroughly after using. Avoid contamination of feed and foodstuffs. Ronnel is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. Atropine is antidote. Consult a veterinarian at first sign of an adverse reaction.

EPA Reg. No. 4476-62

