



SCREW WORM SPRAY

(COLORED BLUE)

KILLS

Screw Worms, Ear Ticks and Blowfly Maggots

Protects Wounds From
Screw Worms and Blowfly Maggots

For livestock of all ages, including newborn animals

ACTIVE INGREDIENTS:


Ronnel [O,O-dimethyl O-(2,4,5-tri-chlorophenyl) phosphorothioate] -----	2.5%
Xylene -----	0.5%
INERT INGREDIENTS -----	97.0%

FOR EXTERNAL USE ONLY
LIVESTOCK REMEDY - NOT FOR HUMAN USE

CAUTION: Keep Out of Reach of Children.
See back panel for additional precautions.

NET WEIGHT 10 OUNCES

MORTON PHARMACEUTICALS, INC.
ETHICAL DISTRIBUTORS — MEMPHIS, TENN. 38108



DIRECTIONS FOR USE:

For use on Beef Cattle, Dairy Cattle, Horses, Sheep, Goats and Swine

Blue color added for marking treated areas

TO OPERATE

Hold outlet about 3 inches from wound or infested area. Be sure opening is directed toward area to be treated. Press valve down to spray, wetting wound and about 1 inch of surrounding area thoroughly.

SCREW WORM AND BLOWFLY INFESTATIONS:

Use particular care to spray into deep pockets caused by screw worms. Repeat applications at 5 to 7 day intervals until wound is healed.

TO PREVENT SCREW WORM AND BLOWFLY INFESTATIONS:

Apply directly to all open or fresh wounds resulting from dehorning, castration, shear cuts, etc. Also treat navels of newborn animals. Freely bleeding wounds may require several applications. Repeat applications at 5 to 7 day intervals until wounds are healed.

EAR TICKS:

Spray directly into animal's ear. Repeat at 5 to 7 day intervals as needed.

FLEECE WORM INFESTATIONS:

Spray infested areas liberally, working toward the center of infestation. Shearing or removing wool from the infested area is not necessary. Re-treat in 5 to 7 days as needed.

CAUTION:

Irritating to Eyes. Avoid contact with skin, eyes and human eyes. In case of contact with human eyes, flush eyes immediately with plenty of water and get prompt medical attention. Avoid breathing spray mist. Wash hands thoroughly after using. Avoid contamination of feed and bedding.

WARNING:

Contents Under Pressure. Do not puncture container. Store at room temperature (not over 120° F.) away from radiators, stoves, hot water or other heat. Never throw container into fire or incinerator when empty. Ronnel is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides or chemicals. Atropine is antidote. Consult a veterinarian at first sign of an adverse reaction.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 4476-63