



DAIRY^{and} FOOD

PROCESSING PLANT

FLY SPRAY

FOR INDUSTRIAL USE

For the Control of

FLIES

and Other Small Flying Insects

ACCEPTED

ACTIVE INGREDIENTS:

Pyrethrins -----	0.3%
Technical Piperonyl Butoxide* -----	0.5%
N-Octyl Bicycloheptene Dicarboximide --	1.1%
Petroleum Distillates -----	98.0%

*Equivalent to 0.48% (Butylcarbityl)(6-Propylpiperonyl) ether and 0.12% other related compounds.

DIRECTIONS: Use in conventional fogging machine. Close room before fogging, leave closed and unoccupied for at least 30 minutes after treatment. Sweep up and destroy all fallen insects.

Dairy Processing Plants, Food Packaging and Food Processing Plants, Restaurants, Homes

For control of flies, mosquitoes, gnats and flying moths: Fog at rate of one-half to one ounce per 1000 cubic feet. Direct fog toward all parts of room and especially windows or light sources which attract these insects. Repeat as necessary.

WARNING: KEEP OUT OF REACH OF CHILDREN.

FLAMMABLE - DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Avoid inhalation of vapors. Harmful if swallowed. If swallowed call physician immediately. In case of skin contact, wash with soap and water. Use only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using.

Remove birds, pets and fish from the area being treated.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

EPA Reg. No. 4476-68

NET CONTENTS 5 GALLONS

**MORTON PHARMACEUTICALS INC.
Memphis, Tenn. 38108**



DAIRY^{and} FOOD

PROCESSING PLANT

FLY SPRAY

FOR INDUSTRIAL USE

For the Control of

FLIES

and Other Small Flying Insects

ACCEPTED

ACTIVE INGREDIENTS:

Pyrethrins -----	0.3%
Technical Piperonyl Butoxide* -----	0.6%
N-Octyl Bicycloheptene Dicarboximide --	1.1%
Petroleum Distillates -----	98.0%

*Equivalent to 0.48% (Butylcarbityl)(6-Propylpiperonyl) ether and 0.12% other related compounds.

DIRECTIONS: Use in conventional fogging machine. Close room before fogging, leave closed and unoccupied for at least 30 minutes after treatment. Sweep up and destroy all fallen insects.

Dairy Processing Plants, Food Packaging and Food Processing Plants, Restaurants, Homes

For control of flies, mosquitoes, gnats and flying moths: Fog at rate of one-half to one ounce per 1000 cubic feet. Direct fog toward all parts of room and especially windows or light sources which attract these insects. Repeat as necessary.

WARNING: KEEP OUT OF REACH OF CHILDREN.

FLAMMABLE - DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Avoid inhalation of vapors. Harmful if swallowed. If swallowed call physician immediately. In case of skin contact, wash with soap and water. Use only when plant is not in operation. Foods should be removed or covered during treatment. All food processing surfaces should be covered or thoroughly cleaned before using.

Remove birds, pets and fish from the area being treated.

Do not reuse empty drum. Send to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

EPA Reg. No. 4476-68

NET CONTENTS 55 GALLONS

**MORTON PHARMACEUTICALS INC.
Memphis, Tenn. 38108**