

DIRECTIONS FOR USE

PREPARATION OF THE SPRAY Fill the spray tank with half the required amount of water, then add the recommended amount of BEL 2,4-D AMINE with agitation and continue filling the spray tank with water. Use enough water per acre to give uniform coverage. The amount of water required for low volume applications may vary from 5 to 25 gallons per acre. For high volume applications, 100 gallons or more of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

TIME OF APPLICATION. Best results are obtained when BEL 2,4-D AMINE is used on weeds that are young and in a rapid growing condition. Applications of lower rates to susceptible annual weeds will be satisfactory, but for perennial weeds and other conditions where weed kill is difficult, use higher rates. When used as a selective spray in crops, the stage of growth of the crop must be considered. Some woody plants and weeds are hard to kill and repeat applications may be necessary.

SMALL GRAIN CROPS (Wheat, Oats, Barley, Rye): Apply BEL 2,4-D AMINE at the rate of 1 to 1 1/4 pints per acre on small grains when fully tillered or stooled (4 to 8 inches tall) but before head emerges from the boot. Injury will be likely to occur if this product is applied earlier than "tiller" or later than "boot". Do not use on grains undersown with legumes such as alfalfa or clovers, except where some legume injury can be tolerated. Do not apply when grain is headed. DO NOT forage or graze treated grain fields within two weeks after treatment with 2, 4-D. DO NOT feed treated straw to livestock.

AERIAL APPLICATION—To small grains (Wheat, Oats, Barley, and Rye) and Rice: Use BEL 2,4-D AMINE at the rate of 1 to 1 1/4 pints in sufficient water to insure adequate coverage.

FLAX: Apply BEL 2,4-D AMINE at the rate of 1 2 to 2 3 pint per acre as weed growth warrants, providing the flax plants have formed 4 to 5 leaves (2 to 6 inches tall). Do not treat flax after the early bud stage.

RICE: In the South and Southwest apply from 1 1/2 to 2 1/2 pints per acre in 5 to 10 gallons of water 7 to 10 weeks after planting, or when the rice is fully tillered but not yet in the boot. In California: Apply 1 1/2 to 2 pints per acre 55 to 65 days after planting, when the rice is through the water and well established, but not after seed stalks emerge. For difficult weeds and where daily temperatures do not exceed 95°F, 3 pints per acre may be used.

SORGHUM: POST-EMERGENCE — Use spray at the rate of 1 pint per acre on plants 6 to 12 inches high when secondary roots are well established. Do not apply from flowering to dough stage. Use drop nozzle when crop is over 10 inches tall.

BEL 2,4-D AMINE

A Selective Weed Killer

U.S.D.A. Reg. No. 4502-4

4 POUNDS 2,4-D ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS: Dimethylamine 2,4-Dichlorophenoxyacetate 49.6%
2, 4-D Chlorophenoxyacetic Acid Equivalent 11.2%

INERT INGREDIENTS: 50.4%

CAUTION

Keep Out of Reach of Children

GENERAL INFORMATION

BEL 2,4-D AMINE is recommended for controlling Arrowhead, Dandelion, Docks, Henbit, Goldenrod, Lambs-quarters, Mustards, Peppergrass, Pigweed, Plantains, Ragweed, Shepherd's-purse, Thistles, Wild Carrot, Wild Morning-Glory, Wild Radish, and many other 2,4-D susceptible broadleaved weeds in crops. This product also can be used on lawns, golf courses, parks, along highways, vacant lots, and other areas without injury to most grasses.

CAUTION

DO NOT take internally. Avoid contact with skin, eyes or clothing. DO NOT store in or fertilizers, seeds, insecticides, or fungicides. DO NOT use the same spray equipment for applying other materials to plants, as injury may result. DO NOT contaminate irrigation ditches or water used for domestic purposes. Although BEL 2,4-D AMINE is non-volatile, care must be used to prevent spray drift from coming into direct contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants susceptible to 2,4-D. Coarse sprays are less likely to drift. DO NOT graze treated areas within seven days after treatment.

NET CONTENTS

55 GALLONS

BEL CHEMICAL & SUPPLY CO., INC.

SCHRIEVER, LOUISIANA

Apr 23-1970
4502-4

CORN: PR
at the rate of
form coverage
3 to 5 day
DO NOT ap
ENCE—apply
spraying corn
extensions
the base of t
ing to dough

SUGAR CAN
acre as a s
lay-by.

PASTURES:
pastures, ma
per acre of
provide for u
weeds are gr
seeded pastu
apply when g
are usually in
Do not graze
seven days af

LAWNS AND
of BEL 2,4 D
obtain unifor
grasses or to
become well
fured or kill
FOR TREAT

of this produ
to one quart
GENERAL W
sides, and a
2,4-D AMINE
oughly wet a
NOTE: Wh
it may be ne
shake throu

WARRAN
SELLER war
chemical des
poses stated
directions n
assumes the
SELLER mak
including on
of FITNESS
SELLER is ou
specific refer
SELLER's li
the purchase
is made.

U

BEL

Copy 23-117c
4903-4

2,4-D AMINE

A Selective Weed Killer

U.S.D.A. Reg. No. 4502-4

4 POUNDS 2,4-D ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS: Dimethylamine 2,4-Dichlorophenoxyacetate 49.6%
2, 4-Dichlorophenoxyacetic Acid Equivalent 41.2%

INERT INGREDIENTS: 50.4%

CAUTION

Keep Out of Reach of Children

GENERAL INFORMATION

BEL 2,4-D AMINE is recommended for controlling Arrowhead, Dandelion, Docks, Henbit, Goldenrod, Lambs-quarters, Mustards, Peppergrass, Pigweed, Plantains, Ragweed, Shepherd's-purse, Thistles, Wild Carrot, Wild Morning-Glory, Wild Radish, and many other 2,4-D susceptible broadleaved weeds in crops. This product also can be used on lawns, golf courses, parks, along highways, vacant lots, and other areas without injury to most grasses.

CAUTION

DO NOT take internally. Avoid contact with skin, eyes or clothing. DO NOT store near fertilizers, seeds, insecticides, or fungicides. DO NOT use the same spray equipment for applying other materials to plants, as injury may result. DO NOT contaminate irrigation ditches or water used for domestic purposes. Although BEL 2,4-D AMINE is non-volatile, care must be used to prevent spray drift from coming into direct contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants sensitive to 2,4-D. Coarse sprays are less likely to drift. DO NOT graze treated areas by dairy animals within seven days after treatment.

NET CONTENTS

55 GALLONS

BEL CHEMICAL & SUPPLY CO., INC.

SCHRIEVER, LOUISIANA

Continued from Opposite Panel)

CORN: **PRE EMERGENCE**--use BEL 2,4-D AMINE at the rate of 1 to 2 quarts in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting, but before corn emerges. **DO NOT** apply to light sandy soils. **POST-EMERGENCE**--apply at the rate of 1 pint per acre. When spraying corn above 10 inches in height, use nozzle extensions ("corn drops"), directing the spray down the base of the corn plant. Do not apply from tasseling to dough stage.

SUGAR CANE: Apply at the rate of 2 to 4 pints per acre as a spring pre-emergence treatment or after lay-by.

PASTURES: To control many broadleaved weeds in pastures, meadows, and rangelands, use 2 to 3 pints per acre of BEL 2,4-D AMINE in sufficient water to provide for uniform application. Treat pastures when weeds are growing actively. Do not apply to recently seeded pastures until grass is well established. Do not apply when grass is in boot to milk stage. Most legumes are usually injured or killed at the rates recommended. Do not graze treated areas by dairy animals within seven days after treatment.

LAWNS AND GOLF FAIRWAYS: Apply 2 to 3 pints of BEL 2,4-D AMINE per acre in sufficient water to obtain uniform coverage. **DO NOT** apply to bent grasses or to freshly seeded turf until grasses have become well established. Most clovers are usually injured or killed. St. Augustine grass may be injured.

FOR TREATING SMALL AREAS: One tablespoonful of this product in 1 1/2 gallons of water is about equal to one quart in 100 gallons of water.

GENERAL WEED CONTROL: On vacant lots, roadsides, and along fence rows, use 2 to 3 quarts of BEL 2,4-D AMINE in 100 gallons of water per acre. Thoroughly wet all foliage to run-off.

NOTE: When stored at temperatures below freezing, it may be necessary to warm up to at least 45°F. and shake thoroughly before using.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and **BUYER** assumes the risk of any use contrary to such directions. **SELLER** makes no other express or implied warranty, including any other express or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of **SELLER** is authorized to do so except in writing with a specific reference to this warranty. In no event shall **SELLER's** liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Reseed thin spots to insure uniform cover
Raise mowing height to 2" during late spr
Water lawns deeply. Occasional deep water
than frequent light sprinklings.

SPREADER SET

(To be determ

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen	12
Available Phosphoric Acid	2
Potash (Water Soluble)	4

ACTIVE INGREDIENTS:

Dimethyl Ester of Tetrachlorotere

INGREDIENTS INERT AS WEED KILLERS:

TOTAL:

NET WEIGHT 1

INTERNATIONAL MINERALS & CHEM
Skokie, Illinois

DIRECTIONS FOR USE

PREPARATION OF THE SPRAY: Fill the spray tank with half the required amount of water, then add the recommended amount of BEL 2,4-D AMINE with agitation and continue filling the spray tank with water. Use enough water per acre to give uniform coverage. The amount of water required for low volume applications may vary from 5 to 25 gallons per acre. For high volume applications, 100 gallons or more of water will be needed for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

TIME OF APPLICATION: Best results are obtained when BEL 2,4-D AMINE is used on weeds that are young and in a rapid growing condition. Applications of lower rates to susceptible annual weeds will be satisfactory, but for perennial weeds and other conditions where weed kill is difficult, use higher rates. When used as a selective spray in crops, the stage of growth of the crop must be considered. Some woody plants and weeds are hard to kill and repeat applications may be necessary.

SMALL GRAIN CROPS (Wheat, Oats, Barley, Rye): Apply BEL 2,4-D AMINE at the rate of 1 to 4 pints per acre on small grains when fully tillered or stooled (4 to 8 inches tall) but before head emerges from the boot. Injury will be likely to occur if this product is applied earlier than "tiller" or later than "boot". Do not use on grains undersown with legumes such as alfalfa or clovers, except where some legume injury can be tolerated. Do not apply when grain is headed. DO NOT forage or graze treated grain fields within two weeks after treatment with 2, 4-D. DO NOT feed treated straw to livestock.

AERIAL APPLICATION—To small grains (Wheat, Oats, Barley, and Rye) and Rice: Use BEL 2,4-D AMINE at the rate of 1 to 4 pints in sufficient water to insure adequate coverage.

FLAX: Apply BEL 2,4-D AMINE at the rate of 1 2 to 2 3 pint per acre as weed growth warrants, providing the flax plants have formed 4 to 5 leaves (2 to 6 inches tall). Do not treat flax after the early bud stage.

RICE: In the South and Southwest apply from 1 1/2 to 2 1/2 pints per acre in 5 to 10 gallons of water 7 to 10 weeks after planting, or when the rice is fully tillered but not yet in the boot. In California: Apply 1 1/2 to 2 pints per acre 55 to 65 days after planting, when the rice is through the water and well established, but not after seed stalks emerge. For difficult weeds and where daily temperatures do not exceed 95°F., 3 pints per acre may be used.

SORGHUM: POST-EMERGENCE—Use spray at the rate of 1 pint per acre on plants 6 to 12 inches high when secondary roots are well established. Do not apply from flowering to dough stage. Use drop nozzle when crop is over 10 inches tall.

BEL 2,4-D AMINE

A Selective Weed Killer

U.S.D.A. Reg. No. 4502-4

4 POUNDS 2,4-D ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS: Dimethylamine 2,4-Dichlorophenoxyacetate 49.6%
2, 4-Dichlorophenoxyacetic Acid Equivalent 11.2%

INERT INGREDIENTS: 50.4%

CAUTION

Keep Out of Reach of Children

GENERAL INFORMATION

BEL 2,4-D AMINE is recommended for controlling Arrowhead, Dandelion, Docks, Henbit, Goldenrod, Lambs-quarters, Mustards, Peppergrass, Pigweed, Plantains, Ragweed, Shepherd's-purse, Thistles, Wild Carrot, Wild Morning-Glory, Wild Radish, and many other 2,4-D susceptible broadleaved weeds in crops. This product also can be used on lawns, golf courses, parks, along highways, vacant lots, and other areas without injury to most grasses.

CAUTION

DO NOT take internally. Avoid contact with skin, eyes or clothing. DO NOT store near fertilizers, seeds, insecticides, or fungicides. DO NOT use the same spray equipment for applying other materials to plants, as injury may result. DO NOT contaminate irrigation ditches or water used for domestic purposes. Although BEL 2, 4-D AMINE is non-volatile, care must be used to prevent spray drift from coming into direct contact with vegetables, flowers, grapes, fruit trees, ornamentals, cotton, or other desirable plants sensitive to 2, 4-D. Coarse sprays are less likely to drift. DO NOT graze treated areas by dairy animals within seven days after treatment.

NET CONTENTS

55 GALLONS

BEL CHEMICAL & SUPPLY CO., INC.

SCHRIEVER, LOUISIANA

(Continued from Opposite Panel)

CORN: PRE-EMERGENCE—use BEL 2,4-D AMINE at the rate of 1 to 2 quarts in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting, but before corn emerges. DO NOT apply to light sandy soils. **POST-EMERGENCE**—apply at the rate of 1 pint per acre. When spraying corn above 10 inches in height, use nozzle extensions ("corn drops"), directing the spray down the base of the corn plant. Do not apply from tasseling to dough stage.

SUGAR CANE: Apply at the rate of 2 to 4 pints per acre as a spring pre-emergence treatment or after lay-by.

PASTURES: To control many broadleaved weeds in pastures, meadows, and rangelands, use 2 to 3 pints per acre of BEL 2,4-D AMINE in sufficient water to provide for uniform application. Treat pastures when weeds are growing actively. Do not apply to recently seeded pastures until grass is well established. Do not apply when grass is in boot to milk stage. Most legumes are usually injured or killed at the rates recommended. Do not graze treated areas by dairy animals within seven days after treatment.

LAWNS AND GOLF FAIRWAYS: Apply 2 to 3 pints of BEL 2,4-D AMINE per acre in sufficient water to obtain uniform coverage. DO NOT apply to bent grasses or to freshly seeded turf until grasses have become well established. Most clovers are usually injured or killed. St. Augustine grass may be injured.

FOR TREATING SMALL AREAS: One tablespoonful of this product in 1 1/2 gallons of water is about equal to one quart in 100 gallons of water.

GENERAL WEED CONTROL: On vacant lots, roadsides, and along fence rows, use 2 to 3 quarts of BEL 2,4-D AMINE in 100 gallons of water per acre. Thoroughly wet all foliage to run-off.

NOTE: When stored at temperatures below freezing, it may be necessary to warm up to at least 45°F. and shake thoroughly before using.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.