



BENZ-O-CHLOR 4-E

FOR CONTROL OF MITES ON CITRUS

RESTRICTED USE PESTICIDE

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses outlined by the certified applicator's certification.

EPA Reg. No. 9518-2

ACTIVE INGREDIENTS

Ethyl 4,4'-dichlorobenzilate	45.5%
kyrene	43.5%

INERT INGREDIENTS

TOTAL	100.0%
-------	--------

Contains 4 lbs. chlorobenzilate per gallon

KEEP OUT OF REACH OF CHILDREN! WARNING

See side panels for additional specific precautionary statements and directions for use.

GENERAL INFORMATION

Tower Benz-O-Chlor 4-E miteicide is an emulsifiable solution intended for application after dilution with water. The rates given (see right panel) — Directions for use are for dilute sprays (rate per 100 gals. of water unless otherwise stated). In general, application should be made when mites first appear. For severe infestations, or if conditions are favorable for mite buildup, repeated applications will be necessary. Do not use with highly alkaline materials such as lime.

NOTICE

Read the label carefully — Use only as directed.

Use of this material other than in accordance with directions on this label may produce plant injury, excessive residues, or other undesirable results. Tower Chemical Company makes no express or implied warranties other than specified on this label.



TOWER CHEMICAL COMPANY
Montverde Rd. and S.C.L. Railroad
Clermont, Florida 32711

EPA EST. NO. 9518-FL-1

DIRECTIONS FOR USE

(Continued from left panel)

CITRUS (lemons, oranges, lemons, limes, tangelos, tangelines, kumquats, Citrus Fruit Mites). Apply 4 to 10 pt. per 100 gals. of water or 5 pts. per acre in sufficient water to assure thorough coverage when mites first appear. For severe infestations or if conditions are favorable for mite buildup, repeated applications will be necessary.

Flat Mites, Six-spotted Mites, Texas Citrus Mites. Apply 10 pt. per 100 gals. of water or apply 5 pt. per acre in sufficient water to assure thorough coverage.

Citrus Red Mites, Purple Mites. Apply 10 pt. per 100 gals. of water or apply 5 pts. per acre in sufficient water to assure thorough coverage for quick cleanup of this pest. For seasonal control, either repeated applications or use in combinations with other miteicides will be required.

Citrus Bud Mites, Pacific Spider Mites, Yuma Spider Mites. Apply 12 to 14 pt. per 100 gals. of water or apply 5 to 7 1/2 pts. per acre in sufficient water to assure thorough coverage.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use only in Arizona, California, Florida and Texas.

Do not use in dwellings.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

WARNING — FLAMMABLE! Keep away from heat or open flame. Store at temperatures above 32°F.

Pesticide Disposal

Pesticide spray mixture or residue that cannot be used or chemically processed should be buried in a safe place away from water supplies.

Container Disposal

(a) Reseal container and offer for reconditioning or

(b) Triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill or bury in a safe place.

Consult Federal, State, or local disposal authorities for approved alternative procedures.

Read the label carefully — Use only as directed.