SANITIZATION OF BEVERAGE CONTAINERS

Follow local Health Department regulations, which usually suggest use-solution in the range of the lower ppm levels shown in the Dilution Table presented below.

DIRECTIONS FOR DILUTION:

Monarch Low Foam Todine	Titratable Iodine
1/2 oz. per 10 Gal Water	6½ ppm
1 oz. per 10 Gal. Water	$12\frac{1}{2}$ ppm
2 oz. per 10 Gal. Wate:	25 ppm
4 oz. per 10 Gal. Water	50 ppm

WARNING:

Do not allow concentrated Monarch Low-Foam lodine to come into contact with food.

Harmful if swallowed.

Avoid contact with skin and eyes. In case of contact with eyes, flush immediately with water, get medical attention.



MONARCH LOW-FOAM IODOPHOR GERMICIDAL DETERGENT

USDA Reg. 4524-19

ACTIVE INGREDIENT:

Butoxy Polypropoxy Polyethoxy Ethanol-Iodine Complex (providing 1.6% titratable icdine)	12.54%
INERT INGREDIENTS	
FOR SANITATION IN:	00.00%
Beer Bottling, Beer Canning, Soft Drink Bottling, Milk Bottl	ing,
Food Cannina	


CAUTION: Keep out of reach of children - See warnings on side panel.

SEP-KO CHEMICALS, INC.

Minneapolis, N

SC378 Rev. 9-68

4524-19



MONARCH LOW-FOAM IODOPHOR GERMICIDAL DETERGENT

USDA Reg. 4524-19

ACTIVE INGREDIENT:

Butoxy Polypropoxy Polyethoxy Ethanol-Iodine Complex (providing 1.6% titratable iodine) 12.54%

INERT INGREDIENTS 87.46%

FOR SANITATION IN: 100.00%

Beer Bottling, Beer Canning, Soft Drink Bottling, Milk Bottling, Food Canning

CAUTION: Keep out of reach of children — See warnings on side panel.

DIRECTIONS FOR USE

EQUIPMENT SANITIZATION

Preclean equipment with cold or lukewarm water.

Fill with solution 2 oz. Monarch Low Foam lodine to 10 gal. water.

(Provides 25 ppm. titratable iodine.)

Equipment must be adequately drained before contact with food.

TANK AND VAT SANITIZATION

Preclean with water (not hot.)

Add solution 2 oz. Monarch Low Foam lodine to 10 gal. water.

(Provides 25 ppm. titratable iodine.)

After 5 minutes contact time, equipment must be adequately drained before contact with food.

Use dilutions providing not more than 25 ppm of titratable iodine do not require a final potable water rinse, in accordance with Federal Food Additive Regulation 121.2547.

SANITIZATION OF WALLS, FLOORS, CONVEYERS, ETC.

Clean daily with solution 2 oz. Monarch Low Foam lodine to 10 gal. water (25 ppm titratable iodine.)

Clean weekly or at batch run and with solution 4 oz. Monarch Low Foam Iodine to 10 gal. water (50 ppm. titratable iodine) and rinse.

MICALS, INC.

Minneapolis, Minnesota 55421