

DIRECTIONS FOR USE

It is a violation of Federal Law to use the product in a manner inconsistent with its labeling.

DILUTION DIRECTIONS: Mix 1 oz. SUPER KABON into 4 gallons water (200 ppm solution).

In all applications an exposure time of at least two minutes should be maintained. The temperature of the solution should be 75°F or above. The pH of the solution should be 6.0 or higher.

Use dilutions providing up to 200 ppm active quaternary (1 oz. SUPER KABON in 4 gallons water) should be used.

FOR THE FOOD PLANT Sanitizing with SUPER KABON

SUPER KABON sanitizing solutions are stable at temperatures up to boiling.

SANITIZING PREVIOUSLY CLEANED SURFACES WITH SUPER KABON: For linings, pipelines, pumps, heaters, vats, and other hard nonporous surfaces, use 1 ounce SUPER KABON per 4 gallons water (200 ppm).

Remove gross food particles and soil. Pre-clean all surfaces. Rinse with potable water prior to application of sanitizing solution. Apply with mop brush or cloth. Thoroughly wet all surfaces. Treated surfaces must remain wet for one minute, followed by adequate draining. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanized applications, use solution may not be recycled for sanitizing applications.

SUPER KABON FOR USE AS A HAND SANITIZING GEL IN FOOD PLANTS

1. Wash hands thoroughly with soap and water.
2. Rinse hands thoroughly.
3. Dip hands in a solution containing SUPER KABON made at the rate of 1 oz. to 4 gallons of water.

(This solution contains 200 ppm active quaternary ammonium compound.)

SANITIZING SHELL EGGS INTENDED FOR FOOD

To sanitize previously cleaned food-grade eggs in shell egg and egg product processing plants, spray with a solution of 1 oz. SUPER KABON in 4 gallons of warm water (200 ppm quat). The solution should be warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs sanitized with this product shall be subjected to a potable water rinse only if they are broken immediately for use in the manufacture of egg products. Eggs should be reasonably dry before casing or breaking. The solution should not be re-used for sanitizing eggs.

FOR THE DAIRY FARM Sanitizing with SUPER KABON

IN THE MILKHOUSE — Sanitize milking equipment by immersion or recirculation with SUPER KABON solution containing one ounce SUPER KABON to 4 gallons of water (200 ppm).

UDDER WASHING — Approximately one minute before applying milking unit, wash and massage teats with a clean paper towel wetted with a solution of SUPER KABON at 1 oz. to 4 gallons of water at 110°F. Provides 100 ppm solution. Never reuse towels or dip a used towel back into solution. Always use freshly prepared solutions and non-corrosible utensils.

Authorized by USDA for use in federally inspected meat and poultry plants. ®

SUPER KABON is a concentrated sanitizing agent and should be diluted before using. SUPER KABON solutions should not be used with soaps or anionic detergents as such mixtures will impair the antibacterial effectiveness of SUPER KABON.

0001G 1087

In Accordance with FR Notice
Based on Fraud Labeling Date 8-18-87

Monarch®

Super Kabon

REG. U.S. PATENT OFFICE

ACTIVE INGREDIENTS: N-Allyl (50% C., 40% C., 10% C.)

dimethyl benzyl ammonium chloride

INERT INGREDIENTS: Water

EPA Reg. No. 4524-24
EPA Est. No. 4524-MN-1

EPA Reg. No. 4524-24
EPA Est. No. 4524-NY-1

EPA Reg. No. 4524-24
EPA Est. No. 4524-CA-2

KEEP OUT OF REACH OF CHILDREN

DANGER:

See side panel for additional precautionary statements



H.B. Fuller Company
Monarch Division

3900 Jackson Street N.E.
Minneapolis, MN 55421

NET CONTENTS: 1 U.S. GAL.
(3.785 LITERS)

Precautionary Statements: Hazards to Humans and Domestic Animals

KEEP OUT OF REACH OF CHILDREN

DANGER:

CORROSIVE, CAUSES SEVERE EYE & SKIN DAMAGE
DO NOT GET IN EYES, ON SKIN OR CLOTHING. WEAR GOGGLES
OR FACE SHIELD, RUBBER GLOVES WHEN HANDLING.

HARMFUL OR FATAL IF SWALLOWED

PREVENT CONTAMINATION OF FOOD. CONSULT MATERIAL
SAFETY DATA SHEET FOR ADDITIONAL INFORMATION

FOR INDUSTRIAL USE ONLY

Medical personnel familiar with Monarch products are
available 24 hours a day, 7 days a week.

Toll-free medical emergency number

1/800-228-5635

STATEMENT OF PRACTICAL TREATMENT: In case of contact immediately flush
eyes or skin promptly with water for at least 15 minutes. For Eyes, call a physician.
Remove and wash contaminated clothing before reuse. If swallowed, drink promptly
a large quantity of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of
gastric lavage. Measures against circulatory shock, respiratory depression, and
convulsion may be needed.

STORAGE AND DISPOSAL

PROHIBITIONS: DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. OPEN DUMP-
ING IS PROHIBITED.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instruc-
tions must be disposed of according to Federal or approved state procedures under Subtitle C of the
Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Plastic Containers: Triple rinse (or equivalent). Then dispose of in a sanitary land-
fill, or by other approved state and local procedures.
GENERAL: CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE
PROCEDURES.

Handwritten notes: 1/800-228-5635, 1/800-228-5635, 1/800-228-5635

H.B. Fuller Company
Monarch Division
3900 Jackson Street NE.
Minneapolis, MN 55421

Attention: Lois Tellefson

Gentlemen:

Subject: Super Kabon
EPA Registration No. 4524-24
Your Amendment Application Dated January 12, 1987

The statement on the label concerning sanitizing of shell eggs with this product as requiring "a potable water rinse only if they are broken immediately for use in the manufacture of egg products" is correct for this product. See #1 on enclosure.

Also, the amendment referred to above is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under section 3(c)(5) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling change listed below before you release the product for shipment bearing the amended labeling:
 - Delete "Ethanol" from the ingredient statement in the active ingredients and include it with inert ingredients.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

15003: I: Lee: L-15: KEMCO: 7/1/87: 7/14/87: AW: VO: dag: ron: VO: SOLJA

CONCURRENCES

SYMBOL	ORIGINATOR							
SURNAME								
DATE								

OFFICIAL FILE CO:

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,


John H. Lee
Product Manager (31)
Disinfectants Branch
Registration Division (TS-767C)

Enclosures

DIRECTIONS FOR USE

It is a violation of Federal Law to use the product in a manner inconsistent with its labeling.
DILUTION DIRECTIONS: Stir 1 oz. MONARCH SUPER KABON in 4 gallons water (200 ppm solution).
 In all applications an exposure time of at least two minutes should be maintained. The temperature of the solution should be maintained. The temperature of the solution should be 75°F or above. The pH of the solution should be 6.0 or higher.
 Use solutions only for up to 24 hours after active quaternary (1 oz. MONARCH SUPER KABON in 4 gallons water) solution is used.

FOR THE FOOD PLANT Sanitizing with MONARCH SUPER KABON

MONARCH SUPER KABON sanitizing solutions are stable at temperatures up to boiling.
SANITIZING PREVIOUSLY CLEANED SURFACES WITH MONARCH SUPER KABON: For kettles, steamers, pumps, freezers, vats, and other hard nonporous surfaces use 1 ounce MONARCH SUPER KABON per 4 gallons water (200 ppm).
 Remove gross food particles and soil. Pre-clean all surfaces. Rinse with potable water prior to application of sanitizing solution. Apply with mop brush or cloth. Thoroughly wet all surfaces. Treated surfaces must remain wet for one minute followed by adequate draining. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanized applications use solution may not be reused for sanitizing applications.

MONARCH SUPER KABON FOR USE AS A HAND SANITIZING DIP IN FOOD PLANTS

1. Wash hands thoroughly with soap and water.
2. Rinse hands thoroughly.
3. Dip hands in a solution containing MONARCH SUPER KABON made at the rate of 1 oz. of MONARCH SUPER KABON to 4 gallons of water.
 (This solution contains 200 ppm active quaternary ammonium compound.)

SANITIZING SHELL EGGS INTENDED FOR FOOD

To sanitize previously cleaned food grade eggs in shell egg and egg product processing plants: spray with a solution of 1 oz. of MONARCH SUPER KABON in 4 gallons of warm water (200 ppm quat). The solution should be warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs sanitized with this product shall be subjected to a potable water rinse only if they are broken immediately for use in the manufacture of egg products. Eggs should be reasonably dry before casing or breaking. The solution should not be re-used for sanitizing eggs.

FOR THE DAIRY FARM Sanitizing with MONARCH SUPER KABON

IN THE MILKHOUSE: Sanitize milking equipment by immersion or circulation with MONARCH SUPER KABON solution containing one ounce MONARCH SUPER KABON to 4 gallons of water (200 ppm).

UDDER WASHING: Approximately one minute before applying milking unit, wash and massage teats with a clean paper towel wetted with a solution of MONARCH SUPER KABON at 1 oz. to 4 gallons of water at 110°F. Provides 100 ppm solution. Never reuse towels or dip a used towel back into solution. Always use freshly prepared solutions and non-corrosible utensils.

MONARCH SUPER KABON is a concentrated sanitizing agent and should be diluted before using. MONARCH SUPER KABON solutions should not be used with soaps or anionic detergents as such mixtures will impair the anti-bacterial effectiveness of MONARCH SUPER KABON.

Authorized by USDA for use in federally inspected meat and poultry plants.

Monarch

SUPER KABON

10% Solution
QUATERNARY GERMICIDE
 REG. U.S. PATENT OFFICE

ACTIVE INGREDIENTS: N-Allyl (50% C., 40% C., 10% C.)

dimethyl benzyl ammonium chloride

Ethanol

INERT INGREDIENTS

EPA Registration No. 4524-200 EPA Est. No. 4524-MN-1

**KEEP OUT OF REACH OF CHILDREN
 DANGER**

See side panel for additional precautionary statements.



H.B. Fuller Company
 Monarch Division

3900 Jackson Street Northeast
 Minneapolis, Minnesota 55421

NET CONTENT

(3.785)