

1E-1001-511-125117

OK 0048

DIRECTIONS FOR USE — GENERAL CLASSIFICATION
IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DISINFECTION IN HOSPITALS, NURSING HOMES AND OTHER HEALTH CARE INSTITUTIONS. For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces. Apply Quik-San with a cloth, mop or mechanical spray device. When applied with a mechanical spray device surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Add 4 1/2 oz. Quik-San to 4 gallons water (850 ppm active quat). Apply to previously cleaned hard surface with a mop or cloth.

At this use level Quik-San is effective against *Pseudomonas aeruginosa*.

DISINFECTANT IN INSTITUTIONS (i.e. Educational, Penal, Military), Industry, Schools and Homes. For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 3 oz. of Quik-San to 5.5 gallons of water (400 ppm active quat). Apply to previously cleaned hard surface with mop or cloth.

At 3 oz / 5.5 gallon use level Quik-San is effective against *Staphylococcus aureus*, and *Salmonella choleraesuis*.

DISINFECTION OF PRECLEANED BARBER TOOLS. Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2 oz. Quik-San to 1 gallon water (380 ppm active quat).

DISINFECTION OF POULTRY EQUIPMENT, ANIMAL QUARTERS AND KENNELS. Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 4 1/2 oz. Quik-San to 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution. Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry or other animals, or empty equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

DISINFECTION OF FOOD EGGS. To sanitize previously cleaned food-grade eggs in shell egg and egg product processing plants, spray with 1 ounce Quik-San in 4 gallons of water (200 ppm active quat) for 1 minute. The solution should be warmer than the eggs but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs sanitized with this product shall be subjected to a potable water rinse only if they are to be immediately broken for use in the manufacture of egg products. Eggs should be reasonably dry before casing or breaking. The solution should not be reused for sanitizing eggs.

Form 1572-285

FUL-SAN

QUIK-SAN

Disinfectant-Sanitizer-Deodorizer

chloride, and sodium
as a preservative for the pe
aged
PA
450

DIRECTIONS FOR USE

SANITIZING OF FOOD PROCESSING EQUIPMENT AND OTHER HARD SURFACES IN FOOD CONTACT LOCATIONS. For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, etc. For countertops, refrigerated storage and display equipment and other hard surfaces. Wash and rinse all articles thoroughly, then apply a solution of 1 oz. Quik-San in 4 gallons of water. Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush. Dishes, silverware, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz / 4 gallon solution of Quik-San. Quik-San does not require a potable water rinse in accordance with 21 CFR 178.1010. Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solutions should be prepared daily or when use solution becomes visibly dirty. For mechanical application, used solution may not be reused for sanitizing applications.

HEAVY DUTY WASHING WITH QUIK-SAN. Approximately 1/2 hour before applying mangle unit, wash and massage towels with a clean paper towel wetted with a solution of Quik-San at 1 oz / 4 gallons of water at 110°F. Never reuse towels or dip a used towel back into solution. Always use freshly prepared solutions.

Precautionary Statements — Hazards To Humans and Domestic Animals

KEEP OUT OF REACH OF CHILDREN

DANGER: Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

STATEMENT OF PRACTICAL TREATMENT — In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

STORAGE AND DISPOSAL

Prohibitions — Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide or rinsings that cannot be used or chemically processed should be disposed of in accordance with local, state and federal regulations.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose in an incinerator or landfill approved for pesticide containers.

GENERAL: Consult Federal, State or Local Disposal Authorities for approved alternative procedures such as limited open burning.

STORAGE: Store away from heat and flames. Protect from freezing.