

PENNWALT

ACCEPTED  
JUL 13 1970  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 4581-27

# BHC W-12

## WETTABLE POWDER

ACTIVE INGREDIENTS  
Benzene hexachloride 12%  
Inert ingredients 88%  
NET WEIGHT 50 LBS.

*Comments*

Pennwalt BHC W-12 is a wetttable micron-sized powder containing 12% gamma isomer of benzene hexachloride. It is compatible with DDT, arsenate of lead, Malathion, Parathion, Aramite, Kryocide, sulfur, neutral coppers, and many other insecticides and fungicides which are non-alkaline in nature.

**FRUIT TREE INSECTS**—Pear Psylla, Prune Thrips, Southern Green Stink Bug and related insects, Rosy, Woolly, Green Apple Aphid, Green Peach Aphid—and certain other aphids and thrips on apple, pear, plums (fresh prunes), and peach trees. Use BHC W-12 at the rate of 2-3 lb. per 100 gal. water, thoroughly wetting foliage. To avoid possible contamination, do not apply within 60 days of harvest or not after fruit starts to form.

**SOUTHERN PINE BARK BEETLE, IPS BEETLE and TURPENTINE BEETLE**—Use 16 lbs. of BHC W-12 per 50 gallons of water, fuel oil or kerosene. Apply to point of run-off, thoroughly wetting trunks of trees while foliage is pale greenish-yellow and before foliage turns dark brown. Birds feeding on treated areas may be killed.

### CAUTION

#### KEEP OUT OF REACH OF CHILDREN

May be Absorbed Through Skin! May produce irritation of skin and eyes. Avoid prolonged breathing of dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Toxic to fish and wildlife. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

USDA Registration No. 4581-27

50 LBS. NET WEIGHT

AGRICULTURAL CHEMICALS DIVISION - PENNWALT CORPORATION

PHILADELPHIA, PA.

PHILADELPHIA, PA.

HOUSTON, TEXAS

RYAN, TEXAS

FRESNO, CALIF.

MONTGOMERY, ALA.

0262A 02A011M

### NOTE

Benzene hexachloride should not be applied under conditions which might result in a toxic residue, or residual odor or taste remaining on food stuffs or edible portions of treated crops. Do not use at higher concentrations than shown on this label. In all cases mix well with water and maintain agitation during spraying. Do not store near any product which will absorb odors. Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

### NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Pennwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Pennwalt Corporation is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential damages such as loss of profits or values.

MADE IN U.S.A.

5-0362B-02A011M

PRINTED IN U.S.A.

REGISTERED UNDER THE FEDERAL  
 PESTICIDE CONTROL ACT  
 FOR ECONOMIC PURPOSES REGISTERED  
 UNDER NO. 1453-2

BHC W-12 is a stable emulsifiable concentrate of 12% gamma benzene hexachloride. It is compatible with DDT, arsenate of lead, Malathion, Parathion, Azinphos, Kelthane, sulfur, neutral coppers, and many other insecticides and fungicides which are non-alkaline in nature.

**FRUIT TREE INSECTS**--Pear Psylla, Pear Thrips, Southern Green Stink Bug and related insects, Rosy, Woolly, Green Apple Aphid, Green Peach Aphid--and certain other aphids and thrips on apple, pear, plums (fresh prunes), and peach trees. Use BHC W-12 at the rate of 2.3 lbs. per 100 gal. water, thoroughly wetting foliage. To avoid possible contamination, do not apply within 70 days of harvest or not after fruit starts to form.

**SOUTHERN PINE BARK BEETLE, IPS BEETLE AND THE PINNACLED BEETLE**--Use BHC W-12 at 1-2 lb. per 100 gallons of water, fuel oil or kerosene. Apply to point of emergence, thoroughly wetting trunks of trees while foliage is pale green. Heavy infestations of foliage may be killed.

**CAUTION**

**CONTAINS 30% OF CHLORIN**

Keep this and all other pesticides out of the reach of children. Do not use near water. Do not apply to crops which are to be consumed fresh. Do not use on grapes, figs, or peaches. Do not use on plants which are to be used for medicinal purposes. Do not use on plants which are to be used for food. Do not use on plants which are to be used for medicinal purposes. Do not use on plants which are to be used for food.

**NOTE**

Always use BHC W-12 in accordance with the directions on the label. Do not use more than the recommended rate. Do not use on plants which are to be consumed fresh. Do not use on grapes, figs, or peaches. Do not use on plants which are to be used for medicinal purposes. Do not use on plants which are to be used for food.

**WARNING**

Do not use on plants which are to be consumed fresh. Do not use on grapes, figs, or peaches. Do not use on plants which are to be used for medicinal purposes. Do not use on plants which are to be used for food.