



# CAUTION

KEEP OUT OF REACH OF CHILDREN

See Specific Caution Statements on Side A CEPTED

USDA Reg. No. 4581-116

50 LB

NET WEIGHT THE FELERAL INSECTIONS FUNCTICIDE AND RODERTICIDE ACE 1 1 SCONOMIC POISON RECEIVED

**AUG 5** 

1971

SOLD BY

A IER NOK-1581-11 AGCHEM-DECCO DIVISION - PENNWALT

#### SUGGESTED USE

powder and suitable for spray or dust application only by ground equipment or airplane. The usual rate of application is 25-50 pounds per acre. A thorough application should be made, covering the underside of the leaves as well as the top. KRYOCIDE is recommended for control of the insects on the crops listed.

BEANS—Mexican bean beetle, bean leaf beetle, bean leaf roller, corn earworm, cucumber beetles.

CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RADISHES, and CAU-LIFLOWER-Imported cabbage worm, cabbage looper, diamond-back moth caterpillar, yellow striped armyworm, climbing cut worms, corn earworm, flea beetles.

CANTALOUPE, WATERMELON, CUCUMBER and SQUASH—Melon worms, pickleworm, cucumber beetles.

TOMATO --- Hornworm, fruitworm, pinworm, blister beetle.

CITRUS - Citrus cutworm, variegated cutworm, orange dog, fruit tree leaf roller, orange tortrix, garden tortrix, Holcocera, Fuller's rose weevil, and katydid.

APPLES, PEACHES, PEARS, SHADE TREES, and ORNAMENTALS-Flea weevil, coddling moth, gypsy moth caterpillar, Fuller's rose beetle, plum curculio.

CRANBERRY —— Cranberry weevil gypsy moth caterpillar, false army worm.

LETTUCE—Cabbage looper.

GRAPES—Grape leaf folder, Omnivorous leaf roller.

STRAWBERRIES—Strawberry leaf roller.

PEPPERS—Pepper weevil.

# CAUTION

Do not apply to corn as injury to the crop may occur. Do not use KRYOCIDE in combination with lime or compounds containing free lime. Do not apply KRYOCIDE immediately before a harvest. Do not apply to bush and vine fruits after fruit starts to form. Do not apply to cabbage, broccoli or grapes after heads or edible parts start to form. Do not apply to collards, mustard cauliflower or lettuce after the seedling stage or to turnips after seedling stage if tops are to be used for feed.

Remove excess residues on edible portions of fruit and vogetables by washing brushing or other effective means.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Burn or bury containers, stay out of smoke; never re-use.

#### NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred



# CAUTION

KEEP OUT OF REACH OF CHILDREN

See Specific Caution Statements on Side ACCEPTED

USDA Reg. No. 4581-116

AUG 5 1971

50 LB NET WEIGHT

FUNCTICIDE AND RODENTICEM ACT

SOLD BY

# AGCHEM-DECCO DIVISION - PENNWALT CORPORATION

TACOMA WASHINGTON

MONROVIA, CALIF

PHILADELPHIA PA

OAK BROOK ILL

BRYAN, TEXAS

TRESNO, CALIF

MONTGOMERY ALA

as the top. KRYOCIDE is recommended for control of the insects on the crops listed.

BEANS—Mexican bean beetle, bean leaf beetle, bean leaf roller, corn earworm, cucumber beetles.

CABBAGE, COLLARDS, MUSTARD, TURNIPS, BROCCOLI, RADISHES, and CAU-LIFLOWER—Imported cabbage worm, cabbage looper, diamond-back moth caterpillar, yellow striped armyworm, climbing cut worms, corn earworm, flea beetles.

CANTALOUPE, WATERMELON, CUCUMBER and SQUASH—Melon worms, pickle-worm, cucumber beetles.

TOMATO—Hornworm, fruitworm, pinworm, blister beetle.

Inologidi abbiication suosia so messi sossimo

CITRUS - Citrus cutworm, variegated cutworm, orange dog, fruit tree leaf roller, orange tortrix, garden tortrix, Holcocera, Fuller's rose weevil, and katydil.

APPLES, PEACHES, PEARS, SHADE TREES, and ORNAMENTALS—Flea weevil, coddling moth, gypsy moth caterpillar, Fuller's rose beetle, plum curculio.

CRANBERRY -- Cranberry weevil, gypsy moth caterpillar, false army worm.

LETTUCE—Cabbage looper.

GRAPES—Grape leaf folder, Omnivorous leaf roller.

STRAWBERRIES—Strawberry leaf roller.

PEPPERS-Pepper weevil.

### **CAUTION**

Do not apply to corn as injury to the crop may occur. Do not use KRYOCIDE in combination with lime or compounds containing free lime. Do not apply KRYOCIDE immediately before a harvest. Do not apply to bush and vine fruits after fruit starts to form. Do not apply to cabbage, broccoli or grapes after heads or edible parts start to form. Do not apply to collards, mustard, cauliflower or lett to after the seedling stage or to turnips after seedling stage if tops are to be used for food.

Remove excess residues on edible portions of fruit and vegetables by washing brushing or other effective means.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Burn or bury containers, stay out of smoke; never re-use.

#### NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Pennwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Pennwalt Corporation is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values.



KRYOCIDE is a f

(96% Sodium Fluoaluminate)

**ACTIVE INGREDIENT:** 

Sodium Fluoaluminate 96.0%
INERT INGREDIENTS 4.0%

TOTAL 100.0%

Fluorine—not less than 50%