ACCEPTED

Hydrothol 191 Granular EPA Reg. No. 4581-172

# DIRECTIONS FOR USE IN CALLYFORNIL SOLLY

inconsistent with its labeling.

It is a violation of Federal law to the this product in inconsistent with its labeling

inconsistent with its labeling. Fungicide, and its annuale Ass, as amended, for the posticide HYDROTHOL 191 GRANU ARE Selected Formulation and submerse aquatic weed killer for use in the fidelity control Southern National Control Co (Najas guadalupensis), American Pondweed (Potamogeton nodosus), Horned Pondweed (Zannichellia palustris), Sago Pondweed (Potamogeton pectinatus), and Coontail (Ceratophyllum demersum).

# TIME OF APPLICATION

Apply 25 to 60 days after sowing, when aquatic weeds are present. Do not apply before rice emerges above surface of water. Do not apply after rice starts heading. Do not apply HYDROTHOL 191 GRANULAR more than once yearly.

## RATE OF APPLICATION

The ideal average depth of water at time of application is 6 to 8 inches; however, based on depth, use rates as follows:

Depth of Water in Inches	Lbs. HYDROTHOL 191 GRANULAR Per Acre	ppm Acid Equivalent
4 – 6	40	2.2 – 1.5
6 8	50	1.9 - 1.4
8 10	60	1.7 — 1.4

### **APPLICATION EQUIPMENT**

Apply with ground or air equipment designed for uniform distribution of granules. Clean out application equipment after each use.

Necessary approval and/or permits should be obtained if required. Consult State Fish and Game agencies before applying this product.

### CAUTION

Do not release water from flooded fields within 10 days of application. Carefully read regular product label and follow appropriate precautions and directions.

PENNWALT — Agchem Division-Pennwalt Corporaty Tacoma, WA Fresno, CA Bryan, TX Philadelphia, PA Columbus, OH