



**ACCEPTED**  
 4581-188  
 DEC 8 1971  
 UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND ROBOTICIDE ACT  
 FOR ECONOMIC PEST CONTROL  
 REGISTERED UNDER NO. 4581-188  
 TO ATTACHED COMMENTS.

# ZINEB W-75

75% Zineb (Wettable Powder)

**ACTIVE INGREDIENT:**

Zineb (Zinc ethylene bisdithiocarbamate) 75.0%  
 (Total Zinc as Metallic 17.7%)

**INERT INGREDIENTS** ----- 25.0%

**TOTAL** ----- 100.0%

## CAUTION

KEEP OUT OF REACH OF CHILDREN

MAY CAUSE IRRITATION OF NOSE AND THROAT. DO NOT BREATHE OR SPRAY MIST. KEEP AWAY FROM FIRE OR SPARKS.

**IMPORTANT**

**COMPATIBILITY:** This product is compatible with most insecticides, organophosphorus compounds, arsenicals and herbicides.

**NOTICE**

Pennwalt Corporation, 10000 W. 10th Avenue, Denver, Colorado 80202. Pennwalt Corporation, 10000 W. 10th Avenue, Denver, Colorado 80202. Pennwalt Corporation, 10000 W. 10th Avenue, Denver, Colorado 80202.

EPA Registration No. 4581-188

NET WEIGHT 3 LBS.

**AGCHEM-DECCO DIVISION—PENNWALT CORPORATION**

TACOMA, WASHINGTON    MONTROVIA, CALIF.    PHILADELPHIA, PA.  
 OAK BROOK, ILL.    BRYAN, TEXAS    FRESNO, CALIF.    MONTGOMERY, ALA.

## GENERAL

## DIRECTIONS FOR USE

Unless otherwise indicated, use 1/2 to 2 lbs. Zineb W-75 per 100 gallons of water. Start filling the tank with water. Sift in the required amount of product when the agitator paddles are covered. Make sure the filter or recirculating pipe from pump are covered. Fill the tank when the product is thoroughly mixed. In regular spray equipment use 100 to 125 gallons of spray per acre for low growing crops. Use 50 to 200 gallons per acre for heavy tomato and potato vines.

**CERREY:** Early blight (Cercospora and Septoria) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water under average conditions every 3 to 5 days. Continue until the crop is harvested. For late blight, apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water after the first sign of the fungus. Continue until the crop is harvested.

**POTATOES:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**PEAS:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**BEANS:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**PEPPERS:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**SPINACH:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**STRAWBERRIES:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**TOMATOES:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**WATERMELONS:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**ZUCCHINI:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**PEPPERS:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.

**STRAWBERRIES:** Late blight (Phytophthora) — start application in the plant bed at first sign of the fungus. Apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested. For early blight (Septoria), apply 1/2 to 1 lb. Zineb W-75 per 100 gallons of water every 3 to 5 days. Continue until the crop is harvested.