

AQUATHOL AQUATHOL

GRANULAR AQUATIC HERBICIDE

ACTIVE INGREDIENT

Dipotassium salt of endothall* 10.1%

INERT INGREDIENTS 89.9%

TOTAL 100.0%

*7-oxabicyclo [2.2.1]heptane-2,3-dicarboxylic acid equivalent 7.2%

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES, hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.

IF SWALLOWED, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

IF ON SKIN, immediately flush with plenty of water for at least 15 minutes. Remove and wash contaminated clothing before reuse.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

See Side Panel for Additional Precautionary Statements



EPA Registration No. 4581-201

EPA Establishment No. 228 IL 1

40 Lbs. Net Weight

AGCHEM DIVISION — ATOCHEM NORTH AMERICA

Philadelphia, PA 19102

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

DANGER

CORROSIVE CAUSES IRREVERSIBLE EYE DAMAGE HARMFUL IF SWALLOWED ABSORBED THROUGH SKIN OR INHALED AVOID CONTACT WITH SKIN OR CLOTHING DO NOT GET IN EYES WEAR GOGGLES OR FACE SHIELD WHEN HANDLING Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Avoid contact with or drift to other crops or plants as injury may result.

Do not use treated water for irrigation or for agricultural sprays on food crops or for domestic purposes within 7 days of treatment.

Do not use fish from treated water for food or feed within 3 days of treatment.

GENERAL INFORMATION

AQUATHOL GRANULAR is a granular aquatic herbicide for use in ponds and lakes which, under field test conditions has shown to be effective against a broad range of aquatic plants with a margin of safety to fish. Dosage rates indicated for the applications of AQUATHOL GRANULAR are measured in "Parts Per Million" (ppm). 1 ppm as a dosage rate means that there would be 1 part of AQUATHOL GRANULAR's active ingredient in 1,000,000 parts of water. Only 1/2 to 5 ppm are generally required for aquatic weed control, whereas some fish species are tolerant to approximately 100 ppm or over. For best results treat areas of one acre or more and/or margins of at least 100 feet in large bodies of water.

Thoroughly clean application equipment immediately after use.

NOTE: Areas treated with AQUATHOL GRANULAR may be used for swimming 24 hours after treatment.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

HOW TO APPLY:

AQUATHOL GRANULAR is a contact killer; consequently, do not apply before weeds are present. Application as early as possible after weeds are present is recommended to permit use of lower application rates. However, for best results water temperature should be at a minimum of 65°F. If an entire pond is treated at one time, or if the dissolved oxygen level is low at time of application, decay of weeds may remove enough oxygen from the water causing fish to suffocate. Water containing very heavy vegetation should be treated in sections to prevent suffocation of fish. Sections should be treated 5-7 days apart. Carefully measure size and depth of area to be treated and determine proper amount of AQUATHOL GRANULAR to apply from chart. For best results apply on a calm day where there is little wave action.

Scatter AQUATHOL GRANULAR as evenly as possible over treated areas. A cyclone seeder is useful for this purpose.

WEEDS CONTROLLED AND AQUATHOL GRANULAR DOSAGE RATE CHART

Common Name	Latin Name	Entire Pond Or Large Area Treatment	Spot Or Lake Margin Treatment
Bass Weed	Potamogeton amplifolius	2.0-3.0 ppm	3.0-4.0 ppm
Bur Reed	Sparganium spp.	3.0-4.0 ppm	4.0-5.0 ppm
Coontail	Ceratophyllum spp.	1.0-2.0 ppm	2.0-3.0 ppm
M. fo.	Myriophyllum spp.	2.0-3.0 ppm	3.0-4.0 ppm
Pondweed			
Bushy	Najas spp.	0.5-1.5 ppm	2.0-3.0 ppm
Curly Leaf	Potamogeton crispus	0.5-1.5 ppm	2.0-3.0 ppm
Flat Stem	Potamogeton zosterifolius	2.0-3.0 ppm	3.0-4.0 ppm
Floating Leaf	Potamogeton natans	1.0-2.0 ppm	2.0-3.0 ppm
Horned	Zannichia spp.	1.0-2.0 ppm	2.0-3.0 ppm
Sago	Potamogeton pectinatus	1.0-2.0 ppm	2.0-3.0 ppm
	Potamogeton nodosus	2.0-3.0 ppm	3.0-4.0 ppm
	Potamogeton diversifolius	1.0-2.0 ppm	2.0-3.0 ppm
	Potamogeton fuliformis	2.0-3.0 ppm	3.0-4.0 ppm
	Potamogeton pusillus	1.0-2.0 ppm	2.0-3.0 ppm
Water Star Grass	Heteranthera spp.	2.0-3.0 ppm	3.0-4.0 ppm

HOW TO DETERMINE QUANTITY TO BE APPLIED

The following charts indicate the total quantity of material to be applied for certain size areas.

APPROXIMATE POUNDS OF AQUATHOL GRANULAR FOR ONE ACRE* TREATMENT DOSAGE IN POUNDS FOR VARIOUS CONCENTRATIONS IN PPM

DEPTH	0.5 ppm	1.0 ppm	1.5 ppm	2.0 ppm	2.5 ppm	3.0 ppm	4.0 ppm	5.0 ppm
1 Ft. Deep	13 lbs.	27 lbs.	41 lbs.	54 lbs.	67 lbs.	81 lbs.	108 lbs.	135 lbs.
2 Ft. Deep	27 lbs.	54 lbs.	81 lbs.	108 lbs.	134 lbs.	161 lbs.	215 lbs.	269 lbs.
3 Ft. Deep	40 lbs.	81 lbs.	121 lbs.	161 lbs.	202 lbs.	242 lbs.	323 lbs.	403 lbs.
4 Ft. Deep	54 lbs.	108 lbs.	161 lbs.	215 lbs.	269 lbs.	323 lbs.	430 lbs.	538 lbs.
5 Ft. Deep	67 lbs.	134 lbs.	202 lbs.	269 lbs.	336 lbs.	403 lbs.	538 lbs.	673 lbs.
6 Ft. Deep	81 lbs.	161 lbs.	242 lbs.	323 lbs.	403 lbs.	484 lbs.	646 lbs.	807 lbs.

FOR 1000 SQUARE FEET TREATMENT

DOSAGE IN POUNDS FOR VARIOUS CONCENTRATIONS IN PPM

DEPTH	0.5 ppm	1.0 ppm	1.5 ppm	2.0 ppm	2.5 ppm	3.0 ppm	4.0 ppm	5.0 ppm
1 Ft. Deep	0.3 lbs.	0.6 lbs.	0.9 lbs.	1.2 lbs.	1.5 lbs.	1.9 lbs.	2.5 lbs.	3.1 lbs.
2 Ft. Deep	0.6 lbs.	1.2 lbs.	1.9 lbs.	2.5 lbs.	3.1 lbs.	3.7 lbs.	4.9 lbs.	6.2 lbs.
3 Ft. Deep	0.9 lbs.	1.9 lbs.	2.8 lbs.	3.7 lbs.	4.6 lbs.	5.6 lbs.	7.4 lbs.	9.3 lbs.
4 Ft. Deep	1.2 lbs.	2.5 lbs.	3.7 lbs.	4.9 lbs.	6.2 lbs.	7.4 lbs.	9.9 lbs.	12.4 lbs.
5 Ft. Deep	1.5 lbs.	3.1 lbs.	4.6 lbs.	6.2 lbs.	7.7 lbs.	9.3 lbs.	12.4 lbs.	15.4 lbs.
6 Ft. Deep	1.9 lbs.	3.7 lbs.	5.6 lbs.	7.4 lbs.	9.3 lbs.	11.1 lbs.	14.8 lbs.	18.5 lbs.

*One acre equals approximately 208' x 208'

Where the area to be treated is greater than those listed in the charts proceed as follows:

a. Compute the approximate surface acreage.

b. Compute the "acre feet".

c. Multiply a by b to determine the total number of acre feet.

d. Multiply the pounds required at the 1 foot depth under the rate to be used by the number of acre feet to determine total quantity to be used.

In instances where the nuisance to be controlled is an exposed surface problem (i.e., some of the broadleaved pond weeds), it is important to get good contact coverage of the problem.

Necessary approval and/or permits should be obtained in states where required.

HOW TO DETERMINE DOSAGE RATE (Active Ingredient)

AQUATHOL GRANULAR is recommended for the control of the following aquatic weeds at the rates indicated. Since AQUATHOL GRANULAR's active ingredient is water soluble and tends to diffuse from the area treated, select the dosage rate applicable to the area to be treated. Use the lower rate in each range of rates when the growth is young and growing and/or where the weed stand is not heavy. Marginal treatments of large bodies of water require highest rates as indicated.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage Instructions: Store in the original container, preferably in a locked storage area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling sweep up spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal Instructions: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal Instructions: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY AND DISCLAIMER

Attochem North America warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. ATOCHEM MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY IN NO CASE SHALL ATOCHEM OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION OR CUSTOMERS, LABOR COST OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

Attochem and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of Attochem.

5-F-1035-02 D5 (3-90)

Made and Printed in U.S.A.