

elf atochem

ATO

PROPOSED LABEL

AQUATHOL® K

AQUATIC HERBICIDE

ACCEPTED

OCT 26 1995

ACTIVE INGREDIENT

Dipotassium salt of endothal*

INERT INGREDIENTS

TOTAL

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 4581-204

40.3%

59.7%

100.0%

*7-oxabicyclo [2.2.1]heptane-2,3-dicarboxylic acid equivalent 28.6%

Contains per gallon 4.23 lb. dipotassium endothal

(equivalent to 3.0 lbs. endothal acid)

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: Measures against circulatory shock, respiratory depression, and convulsion may be needed.

See Back Panel for Precautionary Statements.

For General Information and Directions for Use refer to Accompanying brochure.

EPA Registration No. 4581-204

EPA Establishment No. 228-IL-1

Net Contents _____ Gallons/_____ Liters

ELF ATOCHEM NORTH AMERICA, INC.
Agchem Division

2000 Market Street, Philadelphia, PA 19103

BEST COPY AVAILABLE EPA REC'D

9-29-95

(2)

2 of 4

GENERAL INFORMATION

AQUATHOL K is a liquid concentrate soluble in water which is effective against a broad range of aquatic plants with a margin of safety to fish.

Dosage rates indicated for the application of AQUATHOL K are measured in "Parts Per Million" (ppm) of dipotassium endothall. Only 0.5 to 5.0 ppm are generally required for aquatic weed control, whereas some fish species are tolerant to approximately 100 ppm or over.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

AQUATIC WEEDS CONTROLLED AND DOSAGE RATE CHARTS

AQUATHOL K is recommended for the control of the following aquatic weeds in irrigation and drainage canals, ponds and lakes at the rates indicated. Since the active ingredient is water soluble and tends to diffuse from the area treated, select the dosage rate applicable to the area to be treated. Use the lower rate in each range of rates where the growth is young and growing and/or where the weed stand is not heavy. Marginal treatments of large bodies of water require higher rates as indicated.

HOW TO APPLY:

AQUATHOL K is a contact killer; consequently, do not apply before weeds are present. Application as early as possible after weeds are present is recommended to permit use of lower application rates. However, for best results water temperature should be at 65°F or above. If an entire pond is treated at one time, or if the dissolved oxygen level is low at time of application, decay of weeds may remove enough oxygen from the water, causing fish to suffocate. Water containing very heavy vegetation should be treated in sections to prevent suffocation of fish. Sections should be treated 5-7 days apart. Carefully measure size and depth of area to be treated and determine amount of AQUATHOL K to apply from chart. For best results apply on a calm day where there is little wave action.

AQUATHOL K should be sprayed on the water or injected below the water surface and should be distributed as evenly as possible. It may be applied as it comes from the container or diluted with water depending on the equipment. Some dilution will give better distribution.

In instances where the nuisance to be controlled is an exposed surface problem (i.e., some of the broad-leaved pond weeds) it is important to get good contact coverage utilizing the highest concentration (least water dilution) compatible with the type of equipment used so that even distribution is achieved.

Necessary approval and/or permits should be obtained in states where required.

Common Name	Latin Name	Entire Pond Or Large Area Treatment	Spot Or Lake Margin Treatment
Bass Weed	Potamogeton amplifolius	2.0-3.0 ppm	3.0-4.0 ppm
Bur Reed	Sparganium spp.	3.0-4.0 ppm	4.0-5.0 ppm
Coontail	Ceratophyllum spp.	1.0-2.0 ppm	2.0-3.0 ppm
Hydrilla	Hydrilla verticillata	2.0-3.0 ppm	3.0-4.0 ppm
Milfoil	Myriophyllum spp.	2.0-3.0 ppm	3.0-4.0 ppm
Pondweed			
Bushy	Najas spp.	0.5-1.5 ppm	2.0-3.0 ppm
Curly-Leaf	Potamogeton crispus	0.5-1.5 ppm	2.0-3.0 ppm
Flat-Stem	Potamogeton zosteriformis	2.0-3.0 ppm	3.0-4.0 ppm
Floating-Leaf	Potamogeton natans	1.0-2.0 ppm	2.0-3.0 ppm
Horned	Zannichellia spp.	1.0-2.0 ppm	2.0-3.0 ppm
Sago	Potamogeton pectinatus	1.0-2.0 ppm	2.0-3.0 ppm
	Potamogeton americanus	2.0-3.0 ppm	3.0-4.0 ppm
	Potamogeton diversifolius	1.0-2.0 ppm	2.0-3.0 ppm
	Potamogeton filiformis	2.0-3.0 ppm	3.0-4.0 ppm
	Potamogeton pusillus	1.0-2.0 ppm	2.0-3.0 ppm
Water Star Grass	Heteranthera spp.	2.0-3.0 ppm	3.0-4.0 ppm

RATE OF APPLICATION—LAKES AND PONDS

The following chart indicates the total quantity of material to be applied.

APPROXIMATE GALLONS OF AQUATHOL K FOR ONE ACRE (208' x 208') TREATMENT

DEPTH	DOSAGE IN GALLONS FOR VARIOUS CONCENTRATIONS IN PPM						
	0.5 ppm	1.0 ppm	1.5 ppm	2.0 ppm	3.0 ppm	4.0 ppm	5.0 ppm
1 ft.	0.3	0.6	1.0	1.3	1.9	2.6	3.2
2 ft.	0.6	1.3	1.9	2.6	3.8	5.1	6.4
4 ft.	1.3	2.6	3.8	5.1	7.7	10.2	12.8
6 ft.	1.9	3.8	5.8	7.6	11.5	15.3	19.2

RATE OF APPLICATION—IRRIGATION AND DRAINAGE CANALS**

The following indicates the total quantity of material to be applied.

GALLONS OF AQUATHOL K REQUIRED TO TREAT 1 MILE BY 1 FOOT DEEP*

PPM	WIDTH OF CANAL IN FEET			
	5	10	15	20
1.0 ppm	0.4	0.75	1.2	1.5
2.0 ppm	0.75	1.5	2.3	3.0
3.0 ppm	1.2	2.3	3.5	4.5
4.0 ppm	1.5	3.0	4.5	6.0
5.0 ppm	2.0	3.8	5.7	7.5

The minimum contact time with weeds for optimum results should be 2 hours.

*For deeper water, adjust rate accordingly.

**Not for this use in California.

WARRANTY AND DISCLAIMER

Elf Atochem North America warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. ELF ATOCHEM MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL ELF ATOCHEM OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST; OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

Elf Atochem and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of Elf Atochem.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(AND DOMESTIC ANIMALS)**

DANGER

Corrosive. Causes irreversible eye damage. May be fatal if swallowed or inhaled. Harmful if absorbed through skin. Do not get in eyes, on skin, or on clothing. Do not breathe spray vapors.

APPLICATORS AND OTHER HANDLERS MUST WEAR:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

**User Safety Recommendations:
Users should:**

Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove protective clothing and equipment immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Avoid contact with or drift to other crops or plants as injury may result. Wash out spray equipment with water after each operation.

Do not use fish from treated areas for food or feed within 3 days of treatment. Do not use water from treated areas for watering livestock, for preparing agricultural sprays for food crops, for irrigation or for domestic purposes within the following periods:

Up to 0.5 ppm dipotassium salt
(0.35 ppm acid equivalent)—7 days after application

Up to 4.25 ppm dipotassium salt
(3.0 ppm acid equivalent)—14 days after application

Up to 5.0 ppm dipotassium salt
(3.5 ppm acid equivalent)—25 days after application

Treated water can be used for sprinkling bent grass immediately.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage Instructions: Store in the original container. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. Storage at temperatures below 32°F may result in the product freezing or crystalizing. Should this occur the product must be warmed to 50°F or higher and thoroughly agitated. In the event of a spillage during handling or storage, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal Instructions: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal Instructions: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.