



ACCEPTED
Apr 16, 1970
 FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR TOXIC AND POISON REGISTER-
 ED UNDER NO. 4581-217

TRICARBAMIX

ACTIVE INGREDIENTS:

Zineb (Zinc ethylene bisdithiocarbamate) ----- 45.0%
 (Total Zinc as metallic 10.6%)

Maneb (Manganese ethylenebisdithiocarbamate) -- 15.0%
 (Manganese equivalent as metallic 3.1%)

Ferbam (Ferric dimethyl dithiocarbamate) ----- 15.0%

INERT INGREDIENTS ----- 25.0%

TOTAL ----- 100.0%

U. S. Patents: 2,504,404; 2,710,822

®TRICARBAMIX is a registered trademark of Penwalt Corporation

CAUTION

See Specific Caution Statements on Side Panel

KEEP OUT OF REACH OF CHILDREN

USDA Registration No. 4581-217

Net Weight

AGRICULTURAL CHEMICALS DIVISION — PENWALT CORPORATION

TACOMA, WASHINGTON PHILADELPHIA, PA.
 OAK BROOK, ILL. BRYAN, TEXAS FRESNO, CALIF. MONTGOMERY, ALA.

GENERAL

Tricarbamix is a spray base which may be used for the control of many fungus plant diseases. It mixes readily with water, suspends well, and adheres to sprayed surfaces with a minimum of run off.

DIRECTIONS FOR USE

Use 1/2 to 2 pounds Tricarbamix per acre in 100 to 125 gallons of water by ground application, or in 10 to 12 gallons of water by air application. Start filling the tank with water. Sift in the required amount of product when the agitator paddles are covered or the jet or recirculating pipe from pump is covered. Fill the tank when the product is thoroughly mixed.

CELERY: Early blight (Cercospora) and late blight (Septoria)—start applications in the plant bed and continue in the field as frequently as required (weekly under average conditions, every 3 to 5 days under severe conditions). Remove residues by stripping, trimming and washing.

TOMATOES: Late blight and early blights, Septoria leaf spot, gray leaf spot (Stemphylium), anthracnose—start applications when fruit of first clusters are well formed; repeat at 7 to 10-day intervals. When weather conditions favor late blight, more frequent applications may be necessary starting when plants are set in the fields.

CUCUMBERS, MELONS, SQUASH: Downy mildew and other leaf diseases (except powdery mildew), anthracnose—start treatment as soon as plants appear above ground or when disease threatens and continue at 7 to 10-day intervals as long as necessary. Under severe conditions, shorten interval to 3 to 5 days. For summer squash, do not apply later than 7 days before harvest.

BEANS (FIELD AND SNAP): Rust, anthracnose; **LIMA BEANS:** anthracnose, downy mildew—start applications when disease first appears and repeat at 4 to 7-day intervals as long as disease threatens. Do not apply to beans later than 7 days before any picking.

Do not feed treated plant parts to meat or dairy animals.

CARROTS: Leaf spot diseases—start application when disease threatens and continue at 7 to 10-day intervals as needed. Do not apply later than 7 days before harvest.

OTHER VEGETABLES: ^{UNSPECIFIED} For plant bed use only—Downy mildew of broccoli, cabbage, and lettuce; anthracnose and Cercospora leaf spot of peppers—use 2 lbs. as a drenching spray starting when seed is planted and continue at 7 to 10-day intervals until transplanting.

COMPATIBILITY: This product is compatible with common insecticides, such as methoxychlor, DDT, organophosphorous compounds, arsenicals and rotenone.

CAUTION

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT, AND SKIN. MAY BE HARMFUL IF INHALED OR SWALLOWED. MAY BE ABSORBED THROUGH SKIN AND THROAT.

Do not breathe dust or spray mist. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks. In case of contact, flush with plenty of water; for eyes, get medical attention.

Toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers. Burn or bury containers; never reuse. Stay out of smoke and fumes.

IMPORTANT

Never allow this product to become wet during storage. This may lead to certain chemical changes which will reduce its effectiveness as a fungicide and create vapors which may be flammable. Keep container closed when not in use.

NOTICE

Penwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Penwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Penwalt Corporation is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages such as loss of profits or values.