



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

MAR 3 - 1994

Gary R. Sandberg
ELF ATOCHEM NORTH AMERICA INC.
AGRICHEMICALS DIV.
2000 MARKET ST, 21ST FLOOR
PHILADELPHIA, PA 19103

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Subject: Label Amendment Submission of 08/10/93 in Response to PR Notice 93-7
EPA Reg. No. 4581-255
MANEB 80

Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

WHAT THIS ACCEPTANCE MEANS:

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

WHAT YOU NEED TO DO NEXT:

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

2017

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs (7505C)
U.S. Environmental Protection Agency
401 M Street, SW
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)
Office of Pesticide Programs
Room 266A, Crystal Mall 2
1921 Jefferson Davis Highway
Arlington, VA 22202

Sincerely,



Jim Tompkins, Deputy Chief
Registration Support Branch
Registration Division (7505W)

Attachment

MANEB 80**FUNGICIDE****ACTIVE INGREDIENT:**

Maneb (Manganese ethylenebisdithiocarbamate) 80.0%
 (Equivalent to manganese metallic 16.5%)

INERT INGREDIENTS: 20.0%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN
CAUTION

Si usted no entiende la etiqueta,
 busque a alguien para que se la
 explique a usted en detalle.

Retain English Translation

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, induce vomiting immediately by giving two glasses of water and sticking finger down throat. Call a physician. Never give anything by mouth to an unconscious person.

IF IN EYES, flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

IF ON SKIN, immediately flush skin with plenty of water. Get medical attention if irritation persists.

See Inside Panel for Additional Precautionary Statements

See following pages for General Information, Directions for Use, and Chemigation Instructions. Do not apply this product through any irrigation system unless these instructions are followed.

EPA Registration No. 4581-255

EPA Establishment No. 33910-HL-01

Net Weight 50 Lbs. (22.68 kg)

BEST AVAILABLE COPY

Sold by:

ELF ATOCHEM NORTH AMERICA, INC.—AGCHEM DIVISION
 Philadelphia, PA

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No. 4581-255

MAR 3 1994

ACCEPTED
 WITH COMMENTS
 In EPA Letter Dated

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act
 as amended, for the pesticide
 registered under EPA Reg. No.

ACCEPTED
 WITH COMMENTS
 In EPA Letter Dated

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)

CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH SKIN, OR INHALED
CAUSES MODERATE EYE IRRITATION. Avoid contact with skin, eyes, or
clothing. Avoid breathing mist.

Personal Protective Equipment

**APPLICATORS AND OTHER HANDLERS MUST
WEAR:**

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear when mixing or loading
- Chemical-resistant apron when cleaning equipment,
mixing or loading

Follow manufacturer's instructions
for cleaning/maintaining PPE.
If no such instructions for washables,
use detergent and hot water. Keep and
wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENT

When handlers use closed systems,
enclosed cabs, or aircraft in a
manner that meets the requirements
listed in the Worker Protection
Standard (WPS) for agricultural
pesticides [40 CFR 170.240 (d)
(4-6)], the handler PPE requirements
may be reduced or modified as
specified in the WPS.

Human flaggers must be in enclosed cabs.

The enclosed cabs must be used in a
manner that meets the requirements
listed in the Worker Protection
Standard (WPS) for agricultural
pesticides [40 CFR 170.240 (d)
(4-6)]. The handler PPE requirements
may be reduced or modified as
specified in the WPS.

User Safety Recommendations:
Users should:

Wash hands before eating, drinking,
chewing gum, using tobacco or
using the toilet.

Remove clothing immediately if
pesticide gets inside. Then
wash thoroughly and put on
clean clothing.

Remove PPE immediately after
handling this product. Wash
the outside of gloves before
removing. As soon as possible,
wash thoroughly and change into
clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift and runoff from treated areas may be
hazardous to aquatic organisms in neighboring areas. Do not apply directly
to water, except as specified for the labeled use on cranberries in areas
where surface water is present or to intertidal areas below the mean high
water mark. Do not contaminate water when disposing of equipment
washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law
to use this product in a manner
inconsistent with its labeling.

Do not apply this product in a
way that will contact workers
or other persons, either directly
or through drift. Only protected
handlers may be in the area during
application. For any requirements
specific to your State or Tribe,
consult the agency responsible for
pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance
with its labeling and with the
Worker Protection Standard, 40 CFR
part 170. This Standard contains
requirements for the protection
of agricultural workers on farms,
forests, nurseries, and green-
houses, and handlers of
agricultural pesticides. It
contains requirements for training,
decontamination, notification,
and emergency assistance. It also
contains specific instructions and
exceptions pertaining to the statement
on this label about personal
protective equipment (PPE),
~~notification requirements~~, and
restricted-entry interval. The
requirements in this box only
apply to uses of this product
that are covered by the Worker
Protection Standard.

Do not enter or allow worker
entry into treated areas
during the restricted entry
interval (REI) of 24
hours.

PPE required for early entry
to treated areas that is
permitted under the Worker
Protection Standard and that
involves contact with anything
that has been treated, such
as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

BEST AVAILABLE COPY

INSTRUCTIONS FOR APPLICATION

MANEB 80 may be used as a spray for the control of many fungus plant diseases. It mixes readily with water, suspends well, and adheres to sprayed surfaces with a minimum of runoff.

MANEB 80 is recommended for control of the diseases on the crops listed below.

SPRAYS:

Place MANEB 80 into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on product labels prior to spray preparation.

FOR GROUND APPLICATIONS use at the rate indicated in sufficient water for thorough coverage, minimum 10 gallons. Increased volume of water may be necessary as foliage density increases.

To prepare mixture, sift in the required amount of product when the agitator paddles or the jet or recirculating pipe from pump are covered. Fill the tank with water when the product is thoroughly mixed. Three to four level teaspoonfuls of MANEB 80 per one gallon of water is equivalent to 1½ to 2 pounds per 100 gallons.

FOR AIR APPLICATION use at the rate indicated in sufficient water for thorough coverage a minimum 3 gallons per acre for field crops and 10 gallons per acre for orchard crops.

When dosage ranges are given, use the higher rate and shorter intervals under severe disease pressure, but do not exceed the maximum rate or apply more frequently than the minimum interval given in the directions for that crop.

If this product is used on a crop, no other product containing a different EBDC active ingredient may be used on the same crop during the same growing season.

VEGETABLE CROP USES

CROPS	DISEASES	RATE LBS. ACRE	DIRECTIONS
Beans (Dry)	Anthracnose Downy Mildew Rust	1.5-2.0	Begin when plants are small. Spray on a 5-7 day interval. Do not apply more than 9.6 lbs. active ingredient (12 lbs. product) per acre per season. Do not apply within 30 days of harvest.
Broccoli Brussels Sprouts Cauliflower	Alternaria Leaf Spot Downy Mildew	1.5-2.0	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12 lbs. product) per acre per season. Do not apply within 7 days of harvest.
Cabbage Kohlrabi Chicory Cabbage (lightweight only)	Alternaria Leaf Spot Downy Mildew	1.5-2.0	Plant beds and direct seeded beds: Apply when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12 lbs. product) per acre per season. Do not apply later than 7 days before harvest.
Chinese Cabbage (Bunch Head) Cauliflower and Head Only	Alternaria Leaf Spot Downy Mildew	1.0-1.5	Plant beds and direct seeded beds: Apply when disease threatens. Spray on a 7-10 day interval. Do not apply more than 7.2 lbs. active ingredient (9 lbs. product) per acre per season. Do not apply later than 7 days before harvest in California and 10 days before harvest in Hawaii.
Sweet Corn Popcorn Sweet Corn used for seed production	Common Rust Hemilethia Blight	1.5	East of the Mississippi (except Arkansas & Louisiana): Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 10 lbs. active ingredient (22.5 lbs. product) per acre per season. Do not feed treated forage to livestock.
		1.5	West of the Mississippi (except Arkansas & Louisiana): Use sufficient water for thorough coverage. Start applications when disease first appears at 3-10 day intervals. Add a recommended surfactant or spreader/sticker if needed for better coverage. Do not apply within 7 days of harvest. Do not apply more than 6.0 lbs. active ingredient (7.5 lbs. product) per acre per season. Do not feed treated forage to livestock.
Cucumbers	Alternaria (Blight) Leaf Spot Angular Leaf Spot Anthracnose Downy Mildew Pythium Fruit Rot	1.5-2.0	Begin when disease threatens or plants begin to run. Spray on a 7-10 day interval. Do not apply more than 12.0 lbs. active ingredient (16 lbs. product) per acre per season. Do not apply within 5 days of harvest.
Eggplant	Anthracnose Septoria Leaf Spot Cladosporium Leaf Spot Early Blight Late Blight Gray Leaf Spot (Phytophthora)	1.5-2.0	Begin at first fruit cluster and repeat at 7-10 day intervals. Do not apply more than 11.2 lbs. active ingredient (14 lbs. product) per acre per crop. Do not apply within 5 days of harvest.
Kale	Alternaria Leaf Spot Downy Mildew	1.5-2.0	Begin when disease threatens and apply on a 7-10 day interval. Do not use more than 3.2 lbs. active ingredient (4.0 lbs. product) per cutting. Do not apply within 10 days of harvest.

VEGETABLE CROP USES continued

CROPS	DISEASES	RATE LBS. ACRE	DIRECTIONS
Lettuce (Head and Leaf) Endive	Downy Mildew	1.5-2.0	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12 lbs. product) per acre per season. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming. Cottolene—Spray on a 7-10 day interval. Do not apply more than 6.4 lbs. active ingredient maximum (8.0 lbs. product) per acre per season. Do not apply within 14 days of harvest.
Melons including cantaloupes, cassia, cantaloupes, honeydews, watermelons	Alternaria Leaf Spot Anthracnose Cercospora Leaf Spot Downy Mildew Gummy Stem Blight	1.0-2.0	Apply as soon as plants begin to run or when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.0 lbs. active ingredient (16 lbs. product) per acre per season. Do not apply within 5 days of harvest.
Onions (Dry Bulb)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	2.0-3.0	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 24 lbs. active ingredient (30 lbs. product) per acre per season. Do not apply to exposed bulbs.
Onions (Green)	Botrytis Leaf Blight Downy Mildew Neck Rot Purple Blotch	2.0-3.0	Begin applications when diseases are first reported in the area at 7-day intervals throughout the season. Do not apply within 7 days of harvest. Do not apply more than 16.0 lbs. active ingredient (21 lbs. product) per acre per season. Do not apply to exposed bulbs.
(Furrow Drain)	Smut	3	Apply as a furrow drizzle at time of planting or in seed. Use 75 to 125 gallons of water per acre. Do not use more than 2.4 lbs. active ingredient (3.0 lbs. product) per 25,000 linear feet of row (18 inch spacing). Do not use in California.
Peppers	Anthracnose Cercospora Leaf Spot (Phytophthora) Phytophthora Blight Ripe Rot	1.5-3.0	East of the Mississippi: Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 14.4 lbs. active ingredient (18 lbs. product) per acre per season. Do not apply within 7 days of harvest. West of the Mississippi: Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 9.6 lbs. active ingredient (12 lbs. product) per acre per season. Do not apply within 7 days of harvest.
Potatoes	Early Blight Late Blight	1.5-2.0	Begin when plants are 2 to 6 inches high. Apply on a 5-10 day interval. Do not apply more than 11.2 lbs. active ingredient (14 lbs. product) per acre per season. Do not apply within 14 days of harvest except for Florida, Connecticut, Maine, Massachusetts, New Hampshire, New York, Pennsylvania, Vermont, Wisconsin where the PVI is 3 days. It is recommended that this product be used with an Integrated Pest Management Program. Use IPI should occur 14 days before harvest.
Potatoes (Seed Pieces)	Fusarium Seed Piece Decay	1.0 lb. per 10 gal.	Seed piece treatment only. Dip whole or cut tubers. Sprout in cool place if held before planting. Do not use treated seed pieces for food, feed, or all purposes.
Pumpkins	Angular Leaf Spot Downy Mildew	1.5-2.0	Begin when disease threatens. Spray on a 7-10 day interval. Do not apply more than 12.0 lbs. active ingredient (16 lbs. product) per acre per season. Do not apply within 5 days of harvest.
Squash (Summer and Winter)	Anthracnose Downy Mildew	1.5-2.0	Begin when plants start to run. Spray on a 7-10 day interval. Do not apply more than 12.0 lbs. active ingredient (16 lbs. product) per acre per season. Do not apply within 5 days of harvest.
Tomatoes (Greenhouse and Field)	Anthracnose Septoria Leaf Spot Cladosporium Leaf Spot Early Blight Late Blight Gray Leaf Spot (Phytophthora)	1.5-3.0	East of the Mississippi: Begin at first fruit cluster and repeat on a 7-10 day interval. Do not apply within 5 days of harvest. Do not apply more than 10.0 lbs. active ingredient (12 lbs. product) per acre per crop. West of the Mississippi: Spray on a 7-10 day interval. Do not apply more than 6.4 lbs. active ingredient (8 lbs. product) per acre per season. Do not apply within 5 days of harvest.

BEST AVAILABLE COPY

FIELD CROPS

CROPS	DISEASES	RATE LBS. ACRE	DIRECTIONS
Sugarbeets	Cercospora Leaf Spot	1.5-2.0	Apply when disease appears. Spray on a 7-10 day interval. Do not apply more than 11.2 lbs. active ingredient (14 lbs. product) per season. Do not apply within 14 days of harvest.

0007

FRUIT AND NUT CROPS

CROPS	DISEASES	RATE LBS./ACRE	DIRECTIONS
FOR POME FRUITS Use either the "Pre-Bloom" or "Extended Application" schedules DO NOT COMBINE OR INTEGRATE THE TWO TREATMENT SCHEDULES			
Apples	Rusts (including Cedar apple) Fruit Leaf Spot Physack Scab Sooty Blotch	6.0 or (1.5 lbs. per 100 gal. in dilute spray)	PRE-BLOOM. Begin applications at 1/4 to 1/2 inch green tip and continue on a 7- to 10-day application schedule through bloom. Do not apply more than 19.2 lbs. active ingredient (24 lbs. product) per acre per year. Do not graze livestock in treated areas.
		3.0 or (.75 lbs. per 100 gal. in dilute spray)	EXTENDED APPLICATION OR TARN III: For implementation of IPM programs, applications based on tree-row volume or for use as a resistance management tool: begin applications at 1/4 to 1/2 inch green tip and continue applications on a 7- to 10-day schedule through the second cover spray or to within 77 days of harvest. Do not apply within 77 days of harvest. Do not apply more than 16.8 lbs. active ingredient (21 lbs. product) per acre per year. Do not graze livestock in treated areas. It is recommended that this product be used in an Integrated Pest Management Program. Add spreader sticker to spray mixture for cover sprays and tank mix with an effective systemic protectant/fungicide for more effective control of diseases.
Almonds	Brown Rot Blossom and Twig Blight Scab Shot-hole Fungus (Leaf Blight)	6.0-8.0 or (1.5-2.0 lbs. per 100 gal. in a dilute spray)	Apply at popcorn, full bloom, and petal fall or every 7 to 10 days if bloom is staggered. Omit petal fall spray if early brown rot is present. Do not apply later than 5 weeks after petal fall or within 145 days of harvest. Do not apply more than 25.6 lbs. active ingredient (32 lbs. product) per acre per season. Do not allow livestock to graze in almond orchards.
Bananas	Sigatoka	2.0-3.0	Apply when leaves first appear and repeat as needed on a 14-21 day interval. Do not apply more than 24 lbs. active ingredient (30 lbs. product) per acre per season. May be applied up to the day of harvest.
Cranberries	Fruit Rot	4.8-6.0	Begin applications at mid-bloom and repeat at 7-10 day interval. Do not apply more than 14.4 lbs. (18 lbs. product) per season. Do not apply within 30 days of harvest.
Grapes	Black Rot Bunch Rot Downy Mildew	1.5-4.0 or (0.375-1.0 lbs. per 100 gal. in a dilute spray)	East of the Rocky Mountains Apply in sufficient water to provide thorough coverage starting when shoots are 1/4 to 1 1/2 inches long and continue at 7-10 day intervals until fruit is set. Do not apply within 60 days of harvest. Do not apply more than 19.2 lbs. active ingredient (24 lbs. product) per acre per season.
		1.5-2.5 or (0.375- 0.63 lbs. per 100 gal. in a dilute spray)	West of the Rocky Mountains Apply when shoots are 1/4 to 1 1/2 inches long and continue at 7-10 day intervals. Do not apply more than 6.0 lbs. active ingredient (7.5 lbs. product) per acre per season. Do not apply within 60 days of harvest except in California. California— Do not apply after bloom.
Kodons figs	Surface Mold (Cladosporium) Surface Rot (Alternaria)	0.75 lbs./ 100 gal.	Apply when disease threatens in 100-400 gallons of water acre. Do not apply more than 2.4 lbs. active ingredient (3.0 lbs. product) per acre per season. Do not apply within 10 days of harvest.
Papayas	Anthracnose Phytophthora Fruit Rot Cercospora Black Spot	2.0-2.5 or (0.5- 0.625 lbs. per 100 gal. in a dilute spray)	Apply when disease first threatens and 14-21 day intervals. Do not apply more than 20 lbs. active ingredient per season (26 lbs. product). May be applied up to the day of harvest.

BEST AVAILABLE COPY

7247

**COMMERCIAL ORNAMENTAL CROP USES
(NOT FOR HOME GARDEN USE)
DO NOT USE ANY OF THESE CROPS FOR FOOD OR FEED PURPOSES**

CROP	DISEASES	RATE LBS/100 GAL	DIRECTIONS
Carnations	Alternaria Leaf Spot Anthracnose Botrytis Blight	1.0-1.5	Begin when new growth starts. Repeat weekly.
Chrysanthemums	Ascochyta Ray Blight Botrytis Petiole Spot Septoria Leaf Spot	0.5 lbs or 227 grams	Apply twice weekly during the blooming season for Petiole Spot and Ray Blight. For Septoria Leaf Spot apply weekly throughout the season.
Dahlias	Alternaria Leaf Spot Botrytis Blight	1.0-1.5	Begin when new growth starts. Repeat weekly.
Dogwood	Anthracnose	1.0-1.5	Begin when buds open. Repeat when bracts fall 4 weeks later and in later summer.
Gladiolus	Botrytis Blight Curvatures and Stemphylium Leaf Spots	1.5	Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.
Grasses Do not use on pasture or range grasses. Do not feed clippings to livestock. Do not use if grass is grown for seed.	Brown Patch	0.2 lbs or 90 grams	Begin at first sign of disease. Repeat at 7- to 14-day intervals.
	Dollar Spot	0.4-0.5 lbs or 178-227 grams	Apply as for Brown Patch, above.
	Melting-Out	0.2-0.3 lbs or 85-113 grams	Apply as for Brown Patch, above.
	Leaf, Stem and Stripe Rusts	3.0	Begin when rust pustules are first seen. Repeat at 7- to 14-day intervals. Do not graze treated areas. Do not feed clippings to livestock.
Lilies	Botrytis Blight	1.0-1.5	Begin with new growth. Repeat weekly.
Panacea	Anthracnose	1.0-1.5	Begin with new growth. Repeat weekly.
Poinsettias	Alternaria Leaf Spot Botrytis Blight Phytophthora Blight	1.0-1.5	Apply to foliage and soil in early spring and early fall and 7- to 10-day intervals during the growing season.
Roses	Black Spot Cane Cankers Leaf Spot (Texas) (California)	1.0-1.5	Begin when first leaves unfold. Repeat at 7- to 10-day intervals.
Snapdragons	Rust	1.0-1.5	Begin with emergence. Repeat weekly.
Zinnias	Alternaria Blight Leaf Spot Botrytis Blight	1.0-1.5	Begin with emergence. Repeat weekly.

**CHEMIGATION
NOT FOR THIS USE IN CALIFORNIA
GENERAL INSTRUCTIONS**

Apply this product only through sprinker irrigation systems including center pivot, lateral move, and furrow, side (wheel) roll, traveler, big gun, solid set, or hand move. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect chemigation system to any public water system. Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

SYSTEM REQUIREMENTS

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

INSTRUCTIONS FOR SPRINKLER (OVERHEAD) IRRIGATION:

Observe the requirements in the System Requirements section above.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Apply MANEB 80 only through systems containing anti-siphon and check valves designed to prevent water source contamination or overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shut-off.

Maintain a gentle continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.

Application of more than recommended quantities of irrigation water per acre may result in decreased product performance.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained. In a center pivot system, block the nozzle set nearest the well/pivot/injection unit to prevent spray being applied to this area.

Where sprinkler distribution patterns do not overlap sufficiently, unacceptable disease control may result.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

MANEB 80 may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

Check local restrictions and requirements regarding sprinkler irrigation applications, as they may vary from state to state.

SPRAY PREPARATION:

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

Prepare a suspension of MANEB 80 in a mix tank. Fill the tank with 1/4 or 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of MANEB 80 and then the remaining volume of water.

APPLICATION INSTRUCTIONS:

Set sprinkler system to deliver 0.1 to 1.25 inches of water per acre. Volumes of water higher than this may reduce efficacy. Start sprinkler and then uniformly inject the suspension of MANEB 80 into the irrigation water line so as to deliver the desired rate per acre. The suspension of MANEB 80 should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing.

NOTE: When treatment with MANEB 80 has been completed, do not irrigate the treated area for 24 to 48 hours to prevent washing the chemical off the crop.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage Instructions: Store in the original container in a dry area. If allowed to become wet the product will deteriorate and represent a fire hazard. Keep away from sources of ignition (e.g. sparks and open flame). Close bag when not in use. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, sweep up spillage and dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal Instructions: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal Instructions: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**BEST AVAILABLE COPY
WARRANTY AND DISCLAIMER**

El Atchem North America warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. EL ATACHEM MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. IN NO CASE SHALL EL ATACHEM OR SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT INCLUDING, BUT NOT LIMITED TO, LOSS OF PROFITS, BUSINESS REPUTATION, OR CUSTOMERS; LABOR COST; OR OTHER EXPENSES INCURRED IN PLANTING OR HARVESTING.

El Atchem and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be varied only by agreement in writing signed by a duly authorized representative of El Atchem.

Made and Printed in Holland

5-N311S-02 C2 (3/82)

(8/93)