ACCEPTED LIPATION THE FEDERAL INSECTIONS PERCECUE A D ROUGHTHURE ACT FOR ECONOMIC POISON REGISTERS UNDER NO. 1 (21-01)

TRIMANZONE

ACTIVE INGREDIENTS:

ACTIVE INCREDIENTS.	
Maneb (Manganese ethylenebisdithiocarbamate) (Manganese equivalent às metallic 12%)	00.0%
Zineb (Zinc ethylene bisdithiocarbamate) (Total Zinc as metallic 2.4%)	10.0%
Ferbam (Ferric dimethyl dithiocarbamate)	10.0%
INERT INGREDIENTS	_ 20.0%
TOTAL	100.0%

CAUTION

See Specific Caution Statements on Side Panel

KEEP OUT OF REACH OF CHILDREN

USDA Registration No. 4581-262

Net Weight

SOLD BY

COCCUTUDAL CHEMICALS DIVISION - PERIMALT CORPORATION

TACCHA WASHINGTON

S FERCOF ILL BRYAN, TEXAS

PHILADELPHIA. PA.

FRESNO, CALIF.

MONTGOMERY, ALA.

GENERAL

Trimanzone is a spray base which may be used for the control of many tungus plant diseases. It mixes readily with water, suspends well, and adheres to sprayed surfaces with a minimum of run off.

DIRECTIONS FOR USE

Use 11/2 to 2 pounds Trimanzona per acre in 100 to 125 gallons of water by ground application, or in 10 to 12 gallons of water by air application. Start filling the tank with water. Sift in the required amount of product when the agitator paddies are covered or the jet or recirculating pipe from pump is covered. Fill the tank when the product is thoroughly mixed.

CELERY: Early blight (Cercospora) and late blight (Septoria)—start applications in the plant bed and continue in the field as frequently as required (weekly under average conditions, every 3 to 5 days under severe conditions). Remove residues by stripping irimming and washing.

romatoes: Late and early blights. Septoria leaf spot, gray leaf spot (Stemphylium), anthracnose-start applications when fruit of first clusters are well formed; repeat at 7 to 10-day intervals. When weather conditions favor late blight, more frequent applications may be necessary starting when plants are set in the fields.

CUCUMBERS, MELONS,

SOUASH- Downy mildew and other leaf diseases (except powdery mildew), anthracnose—start treatment as soon as plants appear above ground or when disease threatens and continue at 7 to 10-day intervals as long as necessary. Under severe conditions, shorten interval to 3 to 5 days. For summer squash, do not apply later than 7 days before harvest.

BEANS (FIELD AND SNAP): Rust, anthracnose; LIMA BEANS: anthracnose, downy mildew—start applications when disease first appears and repeat at 4 to 7-day intervals as long as disease threatens. Do not apply to beans later than 7 days before any picking.

Do not feed treated plant parts to meat or dairy animals.

CARROTS: Leaf spot diseases—start application when disease threatens and continue at 7 to 10-day intervals as needed. Do not apply later than 7 days before harvest.

OTHER VEGETABLES: For plant bed use only—Downy mildew of broccoli, cabbage, and lettuce: anthracnose and Cercospora leaf spot of peopers—use 2 lbs. as a drenching spray starting when seed is planted and continue at 7 to 10-day intervals until transplanting.

COMPATIBILITY: This product is compatible with common insecticides, such as methoxychlor, DDT, organophosphorous compounds arsenicals and rotenone.

CAUTION

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT, AND SKIN, MAY BE HARMFUL IF INHALED OR SWALLOWED. MAY BE ABSORBED THROUGH SKIN AND THROAT.

Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or sparks. In case of contact, flush with plenty of water; for eyes, get medical attention.

Toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of wastes and containers.

IMPORTANT

Never allow this product to become wet during storage. This may lead to certain chemical changes which will reduce its effectiveness as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Burn or bury containers, stay out of smoke; never re-use.

NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Pennwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Pennwalt Corporation is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages such as loss of profits or values.

5-9136 -02

Cameraman's Note
Poor Copy