

ACCEPTED
JUN 11 1958
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 4581-263 SUBJECT
TO ALL OTHER COMMENTS.

Net Weight

AGRICULTURAL TECHNICAL

For Manufacturer Use Only

ACTIVE INGREDIENTS:

Dimethyldiuronphosphoramide

INERT INGREDIENTS:

Total

See Antidote statement and other required warning statements on the side panel

KEEP OUT OF REACH OF CHILDREN

USDA Registration No. 4581-



AGRICULTURAL CHEMICALS DIVISION

PENNSALT CHEMICALS CORPORATION

WASHINGTON

PHILADELPHIA, PA.

BRYAN, TEXAS

IRVING, CALIF.

MONTGOMERY, ALA.

DANGER

POISONOUS BY SKIN CONTACT, INHALATION, OR SWALLOWING
RAPIDLY ABSORBED THROUGH SKIN
REPEATED EXPOSURES MAY, WITHOUT SYMPTOMS, BE INCREASINGLY HAZARDOUS
DO NOT GET ON SKIN OR CLOTHING OR IN EYES

Wear clean, heavy rubber gloves, cap, and clean, water-proof or freshly laundered protective clothing (coveralls, etc.) and goggles. Destroy and replace gloves frequently. Wash thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of skin contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

DO NOT BREATHE VAPOR OR SPRAY MIST
Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for dimethyldiuronphosphoramide protection. If handled indoors, provide mechanical exhaust ventilation. Keep all unprotected persons out of the operational area or vicinity.

DO NOT CONTAMINATE FOOD OR FEED PRODUCTS
Bury spillage, clean up area with strong lye solution. Bury containers, never re-use.
Highly toxic to wildlife.

NOTE: Birds feeding in treated areas may be harmed.

ATROPINE IS AN ANTIDOTE
Consult your physician for an emergency supply

FIRST AID TREATMENT

Call a physician at once in all cases of suspected poisoning

If symptoms or signs of poisoning include blurred vision, abdominal cramps and tightness in chest, don't wait for a doctor but give two atropine tablets (each 1/100 gr.) at once.
Remove patient immediately from the area. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces. If swallowed, induce vomiting by giving warm salty or soapy water. Repeat until vomit fluid is clear. Have victim lie down and keep quiet.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON

PHYSICIANS NOTE: Warning symptoms include weakness, headache, tightness in the chest, blurred vision, unreactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Give atropine, preferably by injection, grains 1/100 two or three tablets or more orally or rectally every hour as required up to 10 tablets or until pupils dilate. Never give atropine alone. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

Do not use your spill or storage area as a disposal place.

NOTICE

