

WEED BLITZ™ D

WEED BLITZ D is specifically designed for quick knock-down and control of general mixed vegetation and such perennial grasses as Johnson grass, Dallis grass and reed grass (Phragmites) on noncropped land and industrial sites. WEED BLITZ D may provide season long control of seedlings on treated areas. Soil type, rainfall and length of growing season will influence the residual control of seedlings.

DIRECTIONS FOR USE

APPLICATION: For all uses, uniform application is essential.

WEED BLITZ D may be applied either as a dry granular material or as a water spray. Dissolve in water at rates up to 4 pounds per gallon. Both spray and dry applications become effective as soil treatments following rainfall or irrigation.

For DRY application, conventional herbicide spreaders may be used.

For spray application use equipment with mechanical agitation. Add up to 4 pounds WEED BLITZ D per gallon of water, depending on job requirements, to spray tank. Continue agitation until the herbicide is completely dissolved and during spraying operation. To minimize fire hazard due to dry vegetation, treatment while weeds are small is suggested.

GENERAL VEGETATION CONTROL

WEED BLITZ D will give quick knock-down of vegetation and residual control of broadleaved weeds and grasses on parking lots, fence lines, highway and railroad sites, storage yards and other similar areas. For spray application, add 1/2 to 2 pounds of WEED BLITZ D to sufficient water to cover 100 square feet and apply as a contact spray thoroughly wetting all parts of the plants, particularly the growing tips—1/2 to 1 gallon per 100 square feet is usually adequate for young growth up to 6 inches high. The higher rates may be necessary to insure thorough coverage of heavy vegetation or to provide extended residual control. For DRY granular treatment, apply 1/2 to 2 pounds per 100 square feet. The lighter rates should be used only at time of weed emergence on moist soil or when precipitation is anticipated while weeds are still young. Persistent perennial weeds are most effectively treated when nearing maturity or during dormant stages in the fall. Spring applications may also be made under moderate rainfall conditions. Summer applications are recommended only where rainfall is dependable for leaching the herbicide into the soil. Apply 2 to 4 pounds per 100 square feet in the dry granular form or as a spray. On deep-rooted perennials such as bindweed, Canada thistle, Bermuda grass and quack grass the heavier rates may be required.

NET WEIGHT

A Granular Non-Selective WEED AND GRASS KILLER FOR CONTROL OF GENERAL VEGETATION AND PERENNIAL GRASSES

ACTIVE INGREDIENTS:

Sodium Metaborate Tetrahydrate
($\text{Na}_2\text{B}_2\text{O}_4 \cdot 4\text{H}_2\text{O}$) 66.50%
[Boron Trioxide (B_2O_3) Equivalent 22.7%]

Sodium Chlorate (NaClO_3) 30.00%

Diuron [3-(3,4-dichlorophenyl)-1,
1-dimethylurea] 1.25%

INERT INGREDIENTS: 2.25%

TOTAL 100.00%

U. S. Patent No. 3,032,405

CAUTION

KEEP OUT OF REACH OF CHILDREN

Do not leave container where children or animals may take internally. Harmful if swallowed. Avoid contact with eyes, skin and clothing. May cause irritation of eyes, skin, nose and throat. Avoid contamination of feed and foodstuffs. Do not store in direct contact with ground or concrete floors.

USDA Registration No. 4581-266

SOLD BY

 **PENNWALT**
AGRICULTURAL CHEMICALS

TACOMA, WASH. PHILADELPHIA, PA.
OAK BROOK, ILL. BRYAN, TEXAS FRESNO, CALIF.
MONTGOMERY, ALA.

JOHNSON GRASS CONTROL

WEED BLITZ D is particularly effective on deep-rooted perennial grasses such as Johnson grass and Dallis grass on industrial sites, roadsides, fence lines and storage yards. Apply dry at 1 to 3 pounds per 100 square feet or add 1 to 3 pounds to 1 or 2 gallons of water to adequately cover 100 square feet. Maximum efficiency on Johnson grass is obtained by treating after heading to late fall. Most efficient and economical control is obtained by treating during this period.

IMPORTANT

This material is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops and other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlaid by roots of adjacent valuable growths. Careless application of this material, or washing by water run-off, to areas where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Do not drain or flush equipment on or near such areas.

Do not contaminate domestic or irrigation water or waters used by fish and wildlife. Wash clothing and equipment after application. Keep animals off treated areas until rainfall has leached the herbicide into the soil. Mechanical equipment should be thoroughly cleaned after use.

Burn or bury containers; never re-use. Stay out of smoke and fumes.

NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Pennwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Pennwalt Corporation is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values.

5-9140-01E92M

MADE IN U.S.A.

PRINTED IN U.S.A.

ACCEPTED

WEED BLITZ™ D

NET WEIGHT

A Granular Non-Selective WEED AND GRASS KILLER FOR CONTROL OF GENERAL VEGETATION AND PERENNIAL GRASSES

ACTIVE INGREDIENTS:

Sodium Metaborate Tetrahydrate ($\text{Na}_2\text{B}_2\text{O}_7 \cdot 4\text{H}_2\text{O}$)	66.50%
[Boron Trioxide (B_2O_3) Equivalent 22.7%]	
Sodium Chlorate (NaClO_3)	30.00%
Diuron [3-(3,4-dichlorophenyl)-1, 1-dimethylurea]	1.25%
INERT INGREDIENTS:	2.25%
TOTAL	100.00%

U. S. Patent No. 3,052,405

CAUTION

KEEP OUT OF REACH OF CHILDREN

Do not leave container where children or animals may take internally. Harmful if swallowed. Avoid contact with eyes, skin and clothing. May cause irritation of eyes, skin, nose and throat. Avoid contamination of feed and foodstuffs. Do not store in direct contact with ground or concrete floors.

USDA Registration No. 4581-266

SOLD BY

PENWALT
AGRICULTURAL CHEMICALS

TACOMA, WASH. PHILADELPHIA, PA.
OAK BROOK, ILL. RYAN, TEXAS FRESNO, CALIF.
MONTGOMERY, ALA.

JOHNSON GRASS CONTROL

WEED BLITZ D is particularly effective on deep-rooted perennial grasses such as Johnson grass and Dallis grass on pasture sites, roadsides, fence lines and storage yards. Apply dry at 1 to 3 pounds per 100 square feet or add 1 to 3 pounds to 1 or 2 gallons of water to adequately cover 100 square feet. Maximum efficiency on Johnson grass is obtained by treating after heading to late fall. Most efficient and economical control is obtained by treating during this period.

BEST DOCUMENT AVAILABLE IMPORTANT

This material is not selective in action and may be toxic to all types of vegetation and may render the entire treated area totally or partially unproductive for one or more years. Care should be taken to confine the use or application to the particular area intended to be treated and to avoid its contact with lawns, trees, shrubs, crops and other desirable plants which are not intended to be destroyed or injured. This includes precautions in treating areas which may be underlain by roots of adjacent valuable growth. Careless application of this material, or washing by water run-off, to areas where desirable plants are growing or which will be used for later planting may result in injury to desirable plants. Do not drain or flush equipment on or near such areas.

Do not contaminate domestic or irrigation water or waters used by fish and wildlife. Wash clothing and equipment after application. Keep animals off treated areas until rainfall has leached the herbicide into the soil. Mechanical equipment should be thoroughly cleaned after use.

Burn or bury containers; never reuse. Stay out of smoke and fumes.

NOTICE

Penwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risk referred to therein. Penwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of Penwalt Corporation is authorized to do so except in writing with specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages such as loss of profits or value.

5 9130 0182M

MADE IN U.S.A.

PRINTED IN U.S.A.

DIRECTIONS FOR USE

APPLICATION: For all uses, uniform application is essential.

WEED BLITZ D may be applied either as a dry granular material or as a water spray. Dissolve in water at rates up to 4 pounds per gallon. Both spray and dry applications become effective as soil treatments following rainfall or irrigation.

For DRY application, conventional herbicide spreaders may be used.

For spray application use equipment with mechanical agitation. Add up to 4 pounds WEED BLITZ D per gallon of water, depending on use requirements, to spray tank. Continue agitation until the herbicide is completely dissolved and during spraying operation. To minimize fire hazard due to dry vegetation, treatment while weeds are moist is suggested.

GENERAL VEGETATION CONTROL

WEED BLITZ D will give quick knock-down of vegetation and residual control of broadleaved weeds and grasses in parking lots, fence lines, highway and railroad sites, storage yards and other similar areas. For spray applications, add 1/2 to 2 pounds of WEED BLITZ D to sufficient water to cover 100 square feet and apply as a contact spray thoroughly wetting all parts of the plants, particularly the growing tips—1/2 to 1 gallon per 100 square feet is usually adequate for young growth up to 6 inches high. The higher rates may be necessary to insure thorough coverage of heavy vegetation or to provide extended residual control. For DRY granular treatment, apply 1/2 to 2 pounds per 100 square feet. The lighter rates should be used only at time of weed emergence on moist soil or when precipitation is anticipated while weeds are still young. Persistent perennial weeds are most effectively treated when nearing maturity or during dormant stages in the fall. Spring applications may also be made under moderate rainfall conditions. Summer applications are recommended only where rainfall is dependable for leaching the herbicide into the soil. Apply 2 to 4 pounds per 100 square feet in the dry granular form or as a spray. On deep-rooted perennials such as birdseed, Canada thistle, Bermuda grass and quack grass the heavier rate may be required.

ACCEPTED
8-21-69

4581-266