

GENERAL INFORMATION

TOPSIN M-70 WP is a wettable powder for use on golf course greens, tees, and fairways of Bentgrasses, Bluegrasses, Bermudagrasses, Fescues, Ryegrasses, St. Augustine grasses or their mixtures. TOPSIN M-70 WP is not phytotoxic to any of the above-mentioned grasses when used in accordance with the label. TOPSIN M-70 WP is to be used for the prevention and control of the below-mentioned diseases (provides both curative and protective action).

DIRECTIONS FOR USE

Apply with hand sprayer, power sprayer, or a hose-on proportioner. Add the recommended amount of TOPSIN M-70 WP to a partially-filled spray tank under agitation. Complete filling the tank; continuous agitation is necessary to insure even suspension of the wettable powder. Use spray mixture the same day it is prepared. Spray uniformly over the area to be treated. Apply after mowing or avoid mowing for twelve hours after application.

Apply recommended amounts in sufficient water to obtain thorough coverage, usually five gallons per 1,000 sq. ft. of turf area. When treating golf greens, always treat aprons and approaches. Use the highest recommended rate under conditions of severe disease pressure.

DOLLAR SPOT (*Sclerotinia homoeocarpa*), COPPER SPOT (*Gloeocercospora sorghi*)—Apply TOPSIN M-70 WP at $\frac{3}{4}$ to $1\frac{1}{2}$ oz. per 1,000 sq. ft. beginning when the disease first appears and continue at 10-14 day intervals through the season or as needed.

LARGE BROWN PATCH (*Rhizoctonia solani*)—Apply TOPSIN M-70 WP at $\frac{3}{4}$ to $1\frac{1}{2}$ oz. per 1,000 sq. ft., beginning when the disease first appears and continue at 10-14 day intervals through the season or as needed. When conditions are unusually favorable for development of disease, reduce interval to 5-7 days.

FUSARIUM BLIGHT (*Fusarium roseum*)—Apply TOPSIN M-70 WP at 4-8 oz. per 1,000 sq. ft. Water into the root zone by irrigating with one inch of water immediately after application. Apply two applications at 10-14 day interval, beginning applications when the disease first appears.

STRIPE SMUT (*Ustilago striiformis*)—Apply TOPSIN M-70 WP at 4-8 oz. per 1,000 sq. ft. Water into the root zone by irrigating with one inch of water immediately after application. Apply two applications at 14-21 day intervals. Begin applications when the disease first appears, either in the spring or the fall.



TOPSIN[®] M

70% WETTABLE POWDER

TURF FUNGICIDE

Ornamental Turf Use Only

ACTIVE INGREDIENT:

Dimethyl 4,4' -o-phenylenebis [3-thioallophanate]

INERT INGREDIENTS:

TOTAL

70%
30%
100%

*Nippon Soda Company's trademark for Dimethyl 4,4' -o-phenylenebis [3-thioallophanate] licensed to Pennwalt Corporation

EPA REGISTRATION NO. 4581-288

U. S. Patent Pending

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Specific Caution Statements on Side Panel

NET WEIGHT 2 LBS.

AGCHEM-DECCO DIVISION -- PENNWALT CORPORATION

MONROVIA, CA.
OAK BROOK, IL.

TACOMA, WA.
BRYAN, TX.

PHILADELPHIA, PA.
MONTGOMERY, AL.

May be hazardous
with eyes,
spray mist,
handling. Do
not contain

Keep children
been washed

Do not graze
pings from

EXTERNAL
soap and water

INTERNAL
or mustard
MEDIATELY

Burn or burn
reuse them

Pennwalt Corporation
chemical description
purposes refer
referred to this
or implied warranty
of FITNESS of
Corporation is
reference to this
this warranty
clude consequences

MADE IN U.S.A.

ON

for use on golf
tgrasses, Blue-
ses, St. Augus-
-70 WP is not
d grasses when
M-70 WP is to
the below-men-
and protective

IE

or a hose-on
unt of TOPSIN
nder agitation.
ation is neces-
table powder.
red. Spray uni-
y after mowing
plication.

water to ob-
s per 1,000 sq.
s, always treat
recommended
ressure.

DOLLAR SPOT (*Sclerotinia homoeocarpa*), COPPER SPOT (*Gloeocercospora sorghi*)—Apply TOPSIN M-70 WP at $\frac{3}{4}$ to $1\frac{1}{2}$ oz. per 1,000 sq. ft. beginning when the disease first appears and continue at 10-14 day intervals through the season or as needed.

LARGE BROWN PATCH (*Rhizoctonia solani*)—Apply TOPSIN M-70 WP at $\frac{3}{4}$ to $1\frac{1}{2}$ oz. per 1,000 sq. ft., beginning when the disease first appears and continue at 10-14 day intervals through the season or as needed. When conditions are unusually favorable for development of disease, reduce interval to 5-7 days.

FUSARIUM BLIGHT (*Fusarium roseum*)—Apply TOPSIN M-70 WP at 4-8 oz. per 1,000 sq. ft. Water into the root zone by irrigating with one inch of water immediately after application. Apply two applications at 10-14 day intervals, beginning applications when the disease first appears.

STRIPE SMUT (*Ustilago striiformis*)—Apply TOPSIN M-70 WP at 4-8 oz. per 1,000 sq. ft. Water into the root zone by irrigating with one inch of water immediately after application. Apply two applications at 14-21 day intervals. Begin applications when the disease first appears, either in the spring or the fall.



TOPSIN^{*} M

70% WETTABLE POWDER

TURF FUNGICIDE

Ornamental Turf Use Only

ACTIVE INGREDIENT:

Dimethyl 4,4' -o-phenylenebis [3-thioallophanate]

70%

INERT INGREDIENTS:

30%

TOTAL

100%

*Nippon Soda Company's trademark for Dimethyl 4,4' -o-phenylenebis [3-thioallophanate] licensed to Pennwalt Corporation

EPA REGISTRATION NO. 4581-288

U. S. Patent Pending

CAUTION

KEEP OUT OF REACH OF CHILDREN

See Specific Caution Statements on Side Panel

NET WEIGHT 2 LBS.

AGCHEM-DECCO DIVISION — PENNWALT CORPORATION

MONROVIA, CA.
OAK BROOK, IL.

TACOMA, WA.
BRYAN, TX.

PHILADELPHIA, PA.
MONTGOMERY, AL.

CAUTION

May be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothes before reuse. Do not contaminate feed or food.

Keep children and pets off treated area until material has been washed into the soil and area is completely dry.

Do not graze animals on treated turf. Do not feed clippings from treated turf to livestock and poultry.

ANTIDOTE

EXTERNAL: If spilled on skin, wash off immediately with soap and water. If in eyes, flush with plenty of water.

INTERNAL: Induce vomiting with a mixture of salt, soap or mustard in warm water. CALL A PHYSICIAN IMMEDIATELY.

Burn or bury empty containers, stay out of smoke; never reuse them.

NOTICE

Pennwalt Corporation warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. Pennwalt Corporation makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of Pennwalt Corporation is authorized to do so except in writing with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential damages such as loss of profits or values.

MADE IN U.S.A.

5-0157-01F350

PRINTED IN U.S.A.